

Control of megakaryocyte-specific gene expression by GATA-1 and FOG-1: role of Ets transcription factors

Xun Wang¹, John D. Crispino²,
Danielle L. Letting¹, Minako Nakazawa³,
Mortimer Poncz^{1,3,4} and Gerd A. Blobel^{1,3}

¹University of Pennsylvania School of Medicine, ³Children's Hospital of Philadelphia, Philadelphia, PA 19104 and ²Ben May Institute for Cancer Research, University of Chicago, Chicago, IL 60637, USA

⁴Corresponding author
e-mail: Poncz@email.chop.edu

The transcription factor GATA-1 and its cofactor FOG-1 are essential for the normal development of erythroid cells and megakaryocytes. FOG-1 can stimulate or inhibit GATA-1 activity depending on cell and promoter context. How the GATA-1–FOG-1 complex controls the expression of distinct sets of gene in megakaryocytes and erythroid cells is not understood. Here, we examine the molecular basis for the megakaryocyte-restricted activation of the α IIb gene. FOG-1 stimulates GATA-1-dependent α IIb gene expression in a manner that requires their direct physical interaction. Transcriptional output by the GATA-1–FOG-1 complex is determined by the hematopoietic Ets protein Fli-1 that binds to an adjacent Ets element. Chromatin immunoprecipitation experiments show that GATA-1, FOG-1 and Fli-1 co-occupy the α IIb promoter *in vivo*. Expression of several additional megakaryocyte-specific genes that bear tandem GATA and Ets elements in their promoters also depends on the physical interaction between GATA-1 and FOG-1. Our studies define a molecular context for transcriptional activation by GATA-1 and FOG-1, and may explain the occurrence of tandem GATA and Ets elements in the promoters of numerous megakaryocyte-expressed genes.

Keywords: Ets/Fli-1/FOG/GATA/megakaryocyte

Introduction

Development of hematopoietic cell lineages is controlled by both tissue-restricted and widely expressed transcription factors. The related erythroid and megakaryocytic cell lineages are derived from a common progenitor cell and express an overlapping set of lineage-restricted transcription factors, including GATA-1 and the GATA-1 cofactor, Friend of GATA-1 (FOG-1). However, how these factors control the expression of distinct sets of genes in different lineages is an unresolved question.

GATA-1 is a zinc finger transcription factor that is expressed in erythroid cells, megakaryocytes, mast cells and eosinophils (Weiss and Orkin, 1995). Functional GATA elements are present in the proximal promoters of virtually all erythroid- and megakaryocyte-restricted genes examined. Gene targeting studies revealed that

GATA-1 is required for the normal maturation of both erythroid and megakaryocytic cells (Pevny *et al.*, 1991, 1995; Fujiwara *et al.*, 1996; Shivdasani *et al.*, 1997). FOG-1 was identified based on its ability to specifically bind the N-terminal zinc finger of GATA-1 (Tsang *et al.*, 1997). The expression pattern of FOG-1 resembles that of GATA-1, with the highest levels observed in erythroid cells and megakaryocytes. Mice lacking FOG-1 display an erythroid differentiation block similar to that observed in GATA-1-deficient mice (Tsang *et al.*, 1998), providing *in vivo* evidence that these factors function in the same transcriptional pathway. While FOG-1 deficiency virtually ablates the development of the megakaryocytic lineage (Tsang *et al.*, 1998), megakaryocytes lacking GATA-1 are increased in number, but do not differentiate normally (Shivdasani *et al.*, 1997; Vyas *et al.*, 1999). Recent studies showed that the FOG-1-interacting transcription factor GATA-2 can partially compensate for the loss of GATA-1, thus explaining the less dramatic phenotype in GATA-1 null megakaryocytes (Chang *et al.*, 2002). The importance of direct physical interaction between GATA-1 and FOG-1 is illustrated by the observation that point mutations in the N-terminal zinc finger of GATA-1 that disrupt FOG-1 binding lead to defective erythropoiesis and megakaryopoiesis (Crispino *et al.*, 1999; Nichols *et al.*, 2000; Chang *et al.*, 2002). In transient transfection assays, FOG-1 can repress or activate GATA-1 activity depending on cell and promoter context. For example, while FOG-1 stimulates GATA-1 activity on the p45 NF-E2 gene promoter, which is active in erythroid cells and megakaryocytes (Tsang *et al.*, 1997), it represses GATA-1 activity on the erythroid-specific EKLF and transferrin receptor II promoters, as well as on a synthetic GATA-1-dependent promoter (Fox *et al.*, 1999; Kawabata *et al.*, 2001). There are also examples where other GATA-dependent promoters are repressed by members of the FOG family that are present in various tissues and diverse organisms (Fossett and Schulz, 2001).

The requirement for GATA-1 and FOG-1 for normal erythroid and megakaryocytic differentiation and the dependence of their function on their direct interaction appear at odds with the observation that FOG-1 represses GATA-1 activity on numerous promoters in transfection-based assays. The studies presented here using the megakaryocyte-restricted α IIb gene promoter reveal a molecular context that specifies synergistic gene activation by GATA-1 and FOG-1.

The α IIb gene, which encodes the α integrin chain of the platelet fibrinogen receptor α IIb/ β 3, has long served as a model for understanding the molecular basis of high level, megakaryocyte-specific gene expression. Transient transfection and transgenic mice studies showed that ~900 bp of α IIb promoter upstream sequence are sufficient to direct megakaryocyte-specific expression of a linked

reporter (Uzan *et al.*, 1991; Prandini *et al.*, 1992; Block *et al.*, 1994; Tronik-Le Roux *et al.*, 1995). Two pairs of GATA and Ets consensus binding motifs in the 5'-flanking region of the α IIb gene contribute to high level, tissue-specific expression (Uzan *et al.*, 1991; Prandini *et al.*, 1992). Neighboring GATA and Ets elements have been identified in numerous megakaryocyte-specific regulatory regions and are viewed as hallmarks of megakaryocyte-expressed genes. Yet, how these elements function in concert is not understood. While GATA-1 can bind both α IIb GATA elements *in vitro* (Romeo *et al.*, 1990) and, besides GATA-2, is probably the major GATA-binding activity in these cells, the nature of the Ets-binding proteins is more complex. Thus, Ets-1, Ets-2 and Fli-1, but not PU.1 (Spi-1), can bind *in vitro* to the Ets site closest to the transcriptional start site (TSS), whereas PU.1, but not Fli-1 or Ets-1, can bind to the distal Ets element (Lemarchandel *et al.*, 1993; Zhang *et al.*, 1993; Doubeikovski *et al.*, 1997). Transient expression studies in non-megakaryocytic cell lines have shown that GATA-1 can stimulate the expression of a reporter gene containing 75 bp of the proximal α IIb promoter (Lemarchandel *et al.*, 1993). More recently, it was shown that stably expressed GATA-1 and FOG-1 cooperated during the activation of the α IIb promoter in heterologous cells (Gaines *et al.*, 2000). However, it remained unresolved whether activation resulted from direct occupation of the α IIb regulatory elements by both GATA-1 and FOG-1, and whether it required physical interaction between them.

To elucidate the basis for megakaryocyte-specific expression of a GATA-1-dependent gene, our studies focused on the regulation of the α IIb promoter. We found that activation of this promoter by GATA-1 and FOG-1 is dependent on their direct physical interaction and requires the presence of a specialized Ets element. This Ets element, when placed next to an isolated GATA site, enabled GATA-1/FOG-1 synergy. The hematopoietic Ets protein Fli-1 binds to this site and stimulates GATA-1/FOG-1-dependent transcription. Finally, *in vivo* expression of several megakaryocyte-restricted genes, whose promoters contain GATA and Ets elements, depends on direct interaction between GATA-1 and FOG-1. These studies suggest that cooperativity between GATA-1, FOG-1 and Fli-1 is important for establishing and/or maintaining the megakaryocytic cell lineage.

Results

Physical interaction between GATA-1 and FOG-1 is required for α IIb promoter activation

To examine the molecular mechanism by which GATA-1 and FOG-1 activate expression of the megakaryocyte-restricted α IIb gene, we performed transient transfection studies using the α IIb promoter fused to the human growth hormone reporter gene. Expression of GATA-1 alone led to 5- and 8-fold activation of the murine and rat α IIb promoters, respectively (Figure 1A), consistent with previous reports (Lemarchandel *et al.*, 1993; Gaines *et al.*, 2000). While expression of FOG-1 alone had little or no effect on reporter activity, co-expression of GATA-1 and FOG-1 activated these promoters 45- (murine) and 30-fold (rat) (Figure 1A). Control western blots showed




Fig. 1. Dependence of α IIb promoter activation on physical interaction between GATA-1 and FOG-1. (A) Activities of mouse (m, 946 bp) and rat (r, 912 bp) α IIb promoters in transiently transfected NIH-3T3 cells. GATA-1 and FOG-1 were co-expressed as indicated. Promoterless human growth hormone (GH) reporter and the synthetic GATA-1-dependent M1 α reporter served as controls. Means \pm SE are shown. Lower panel: anti-GATA-1 western blot. (B) Experiments were performed as in (A) using the mouse α IIb promoter.

that FOG-1 did not alter the expression of GATA-1, indicating that the effects of GATA-1 and FOG-1 are due to functional synergy at the α IIb promoter (Figure 1A). When assayed on a synthetic reporter gene construct that contains a single GATA element (M1 α ; Martin and Orkin, 1990), FOG-1 repressed GATA-1 activity \sim 3-fold, similar to the findings of a previous report (Fox *et al.*, 1999). These results suggest that the α IIb promoter contains a functional element(s) that leads to stimulation of GATA-1 activity by FOG-1 and is absent in M1 α .

To determine whether FOG-1 function is mediated directly through GATA-1 or through other transcription factors bound at the α IIb promoter, we examined whether the observed functional synergy requires direct interaction

between GATA-1 and FOG-1. To this end, we utilized mutant versions of GATA-1 and FOG-1 that alter their association. GATA-1^{V205G} contains a point mutation in the N-terminal zinc finger of GATA-1 that diminishes FOG-1 binding without affecting its ability to bind DNA (Crispino *et al.*, 1999). This mutation impairs the ability of GATA-1 to trigger terminal differentiation of GATA-1-dependent proerythroblasts (Crispino *et al.*, 1999). A mutation at the same residue of GATA-1 was identified in male patients with X-linked dyserythroblastic anemia and thrombocytopenia (Nichols *et al.*, 2000). FOG-1^{S706R} contains a single amino acid substitution in its sixth zinc finger that restores binding to GATA-1^{V205G}. Expression of FOG-1^{S706R} in erythroid cells expressing GATA-1^{V205G} rescues GATA-1-dependent erythroid differentiation (Crispino *et al.*, 1999). Both wild-type GATA-1 and GATA-1^{V205G} alone activated the α Ib promoter with similar efficiencies (Figure 1B). However, FOG-1 stimulated the activity of GATA-1^{V205G} substantially less than wild-type GATA-1, indicating that physical interaction between GATA-1 and FOG-1 is required for their synergy. To rule out the possibility that GATA-1^{V205G} has additional defects unrelated to its impaired FOG-1 interaction, we examined the effects of co-expressed FOG-1^{S706R} on α Ib promoter activity. The results show that FOG-1^{S706R} displayed strong transcriptional synergy with GATA-1^{V205G}, comparable to that seen with their wild-type counterparts (Figure 1B). These findings demonstrate the importance of direct interaction between GATA-1 and FOG-1 during activation of the α Ib gene promoter *in vivo*.

FOG-1 and GATA-1 occupy the proximal α Ib promoter region *in vivo*

The above studies indicate that GATA-1 and FOG-1 regulate the expression of the α Ib gene by binding to its promoter region. To demonstrate that both GATA-1 and FOG-1 occupy this region *in vivo* in an appropriate cellular context, we performed chromatin immunoprecipitation (ChIP) assays, using the murine megakaryocytic cell line Y10, which expresses endogenous α Ib, GATA-1 and FOG-1 (Ishida *et al.*, 1993; data not shown). Antibodies against GATA-1 and FOG-1, but not isotype-matched control antibodies, immunoprecipitated chromatin that was enriched for proximal α Ib promoter sequences (–110 bp to +226 bp) (Figure 2). As a negative control, a more distal α Ib promoter domain, between –3.3 and –3.6 kb upstream of the α Ib TSS which does not contain a GATA-1-binding site, was not enriched. As an additional control, the promoter region of the housekeeping gene GAPDH, which is not regulated by GATA-1 (Tsang *et al.*, 1998), was not bound by GATA-1 or FOG-1 (Figure 2). These results show that GATA-1 and FOG-1 specifically contact the domain of the α Ib promoter that contains a functionally important GATA element, supporting the hypothesis that transcriptional activation of the α Ib gene is mediated directly by a GATA-1–FOG-1 complex *in vivo*.

60 bp of α Ib upstream sequence are sufficient to mediate GATA-1/FOG-1 synergy

To identify the minimal region of the α Ib promoter sufficient for mediating synergistic activation by GATA-1 and FOG-1, we examined a series of 5' to 3' deletion




Fig. 2. Occupancy of the α Ib promoter by GATA-1 and FOG-1 *in vivo*. ChIP assays using the murine megakaryocytic cell line Y10, anti-FOG-1 and anti-GATA-1 antibodies or isotype-matched control antibodies. Purified DNA fragments were amplified by PCR with primers specific for the proximal mouse α Ib promoter (–110 to +226 bp) and, as control, a distal region lacking GATA and Ets elements (–3.3 to –3.6 kb). The GATA-1-independent housekeeping gene GAPDH served as control. One of three independent experiments is shown.




Fig. 3. Sixty base pairs of upstream region are sufficient for GATA-1/FOG-1 synergy. Promoter constructs containing the indicated lengths of upstream region were analyzed as in Figure 1.

constructs in transiently transfected NIH-3T3 cells. The results revealed that a construct spanning from –60 to +32 bp displayed GATA-1/FOG-1 synergy comparable with that observed with the full-length promoter (Figure 3). This region contains functional, phylogenetically conserved GATA and Ets elements, suggesting that these sites might be sufficient for transcriptional synergy by GATA-1 and FOG-1.

Specificity of the Ets element determines transcriptional output by GATA-1 and FOG-1

To determine the *cis*-acting elements that confer activation by GATA-1 and FOG-1, we compared the α Ib promoter with the M1 α promoter, where FOG-1 inhibits GATA-1 activity (Figure 1A; Fox *et al.*, 1999). We noted that the M1 α promoter contains an Ets element positioned at the same distance from the GATA site as that found in the α Ib promoter (Figure 4A). However, the M1 α and α Ib Ets elements differ from each other in the nucleotides flanking the core 5'-GGA-3' sequence (Figure 4A). These nucleotide differences are predicted to alter Ets binding specificity (Graves and Petersen, 1998). To examine whether the α Ib Ets motif determines GATA-1/FOG-1




Fig. 4. The α IIb Ets element can mediate GATA-1/FOG-1 synergy on a synthetic promoter construct. (A) Sequence alignment of α IIb in the 60 bp construct and in the M1 α GATA and Ets elements. (B) Insertion of the α IIb Ets element into M1 α leads to activation of GATA-1 by FOG-1. G, GATA-1-binding site; E, Ets-binding site; U, 5'-UTR. Means \pm SE are shown.

activity, we replaced the M1 α Ets element with that derived from the α IIb gene. Remarkably, the presence of the α IIb Ets element converted FOG-1 from an inhibitor to an activator of GATA-1 activity (Figure 4B). Substitution of the M1 α 5'-untranslated region (5'-UTR) with that derived from α IIb further increased GATA-1 and FOG-1 synergy (Figure 4B). Thus, these data suggest that the α IIb Ets site is sufficient to convert FOG-1 from a GATA-1 repressor into a co-activator, and that additional sequences in the α IIb regulatory region contribute to maximal activation of GATA-1 by FOG-1.

Fli-1 can bind to the α IIb Ets element *in vitro* and *in vivo*

Given the importance of the Ets element during GATA-1- and FOG-1-mediated transcription, we used electrophoretic mobility shift assays (EMSAs) to characterize the protein(s) that bind to the α IIb and M1 α Ets elements. Using nuclear extracts from NIH-3T3 cells, protein complexes with distinct mobilities were observed that showed little or no cross-competition (Figure 5A), suggesting that these elements bind different members of the Ets protein family. When Y10 cell extracts were used for the EMSA, an additional, faster moving complex was detected with the α IIb Ets probe (Figure 5B). This complex was undetectable with NIH-3T3 cell extracts (Figure 5A) or when the M1 α Ets element was used as probe (Figure 5B). This suggests that at least one Ets factor binds in both a site- and a tissue-specific manner.

The Ets family protein Fli-1 is a likely candidate for this Ets-binding activity since recombinant Fli-1 can bind the proximal α IIb Ets element *in vitro*, and this element matches a consensus Fli-1-binding site (Zhang *et al.*, 1993; Mao *et al.*, 1994; Szymczynska and Arrowsmith, 2000).




Fig. 5. Selective binding of Fli-1 to the α IIb Ets element. (A) α IIb and M1 α Ets elements bind distinct Ets proteins. EMSA using nuclear extracts from NIH-3T3 cells. Note that there is little or no cross-competition between the α IIb and M1 α Ets elements. (B) Nuclear extracts from Y10 cells (left panel) and fetal liver-derived primary megakaryocytes (right panel) were used. Anti-Fli-1 (α Fli-1), but not control (ctr), antibodies reacted with a band (thick arrow), resulting in a supershift (thin arrow).

Moreover, Fli-1 is essential for the normal development of the megakaryocytic lineage (Hart *et al.*, 2000; Spyropoulos *et al.*, 2000; Kawada *et al.*, 2001). To test directly whether Fli-1 is the megakaryocyte-specific Ets-binding protein, we added anti-Fli-1 antibodies to the EMSA reaction. The results show that anti-Fli-1 antibodies supershifted the megakaryocyte-specific band, but not the other Ets-binding activities (Figure 5B). In contrast, anti-Ets-1, anti-Ets-2 and anti-PU.1 antibodies did not alter the mobility of this band (data not shown). To determine whether Fli-1 DNA-binding activity is present in primary megakaryocytes, fetal liver cells were expanded in culture in the presence of thrombopoietin (TPO), leading to enrichment of megakaryocytes up to 80% of the total population of cells as determined by acetylcholinesterase (AChE) staining (data not shown). Nuclear extracts from these cells yielded protein complexes similar to those found in Y10 cells, and included a protein that specifically reacted with anti-Fli-1 antibodies (Figure 5B). However, anti-Fli-1 antibodies consistently supershifted only a fraction of this complex, suggesting that other megakaryocyte-expressed Ets family members also bind the α IIb Ets element (Figure 5B).




Fig. 6. Occupancy of the α Ib promoter by Fli-1 *in vivo*. ChIP assays using Y10 cells, rabbit and mouse anti-Fli-1 antibodies or isotype-matched control antibodies. Primer pairs were the same as in Figure 2. The GAPDH gene served as control.

To determine whether Fli-1 occupies the α Ib promoter *in vivo*, we performed ChIP assays, using Y10 cells. Two anti-Fli-1 antibodies, but not isotype-matched control antibodies, immunoprecipitated chromatin that was enriched for proximal α Ib promoter sequences (–110 bp to +226 bp) (Figure 6). In contrast, a more distal domain, between –3.3 and –3.6 kb upstream of the α Ib transcription initiation site, was not enriched. These results show that Fli-1 binds *in vivo* to the proximal α Ib promoter and suggest that Fli-1 is a strong candidate Ets factor that mediates transcriptional synergy by GATA-1 and FOG-1.

Fli-1 mediates GATA-1/FOG-1 synergy at the α Ib promoter

To determine whether Fli-1 can mediate transcriptional synergy by GATA-1 and FOG-1, we performed transient transfection assays but, to avoid interference by endogenous Ets proteins, we used an α Ib reporter gene construct in which the Ets element had been replaced with a GAL4-binding site (pGL2- α Ib100). While GATA-1 alone weakly activated this construct, FOG-1 co-expression failed to augment GATA-1 activity, consistent with a requirement for an Ets protein activity for α Ib gene expression (Figure 7). We then generated a construct in which the N-terminal 274 amino acids of Fli-1, lacking the DNA-binding domain, were fused to the DNA-binding domain of the yeast transcription factor GAL4 (GAL4-Fli-1). When GAL4-Fli-1 was co-expressed together with GATA-1 and FOG-1, transcriptional activation was restored (Figure 7). In contrast, expression of only the GAL4 DNA-binding domain failed to activate this reporter in the presence of GATA-1 and FOG-1. Control western blots showed that neither FOG-1 nor GAL-4 fusion proteins altered GATA-1 expression (data not shown). These results suggest that Fli-1 is sufficient to mediate transcriptional activation by GATA-1 and FOG-1. The effects of Fli-1 were specific since a construct containing only amino acids 1–194, which lacks an important protein interaction domain required for Fli-1 activity (Watson *et al.*, 1997) was ineffective (Figure 7). To assess whether other Ets family proteins can substitute for Fli-1, we examined a construct in which the N-terminal 160 amino acids of the myeloid/lymphoid Ets protein PU.1 were fused to GAL4. The GAL4-PU.1 construct is functional, since it was used previously to activate gene expression in a myeloid-specific fashion (Maitra and Atchison, 2000). In addition, we generated a construct in which GAL4 was




Fig. 7. Fli-1 mediates GATA-1/FOG-1 synergy on the α Ib promoter. Inset: schematic of the assay. The reporter construct contained 100 bp of α Ib promoter upstream region in which a GAL4-binding site was substituted for the Ets element. GAL4 fusion constructs were co-expressed as indicated.

fused to the lymphoid Ets factor Spi-B. Both GAL4-PU.1 and GAL4-Spi-B failed to activate the α Ib promoter in the presence of GATA-1 and FOG-1, indicating that the effects of Fli-1 are specific (Figure 7; data not shown).

Interaction between GATA-1 and FOG-1 is required for the expression of multiple megakaryocyte-specific genes *in vivo*

The frequent presence of tandem GATA- and Ets-binding sites in the regulatory regions of megakaryocyte-expressed genes has been noted previously (reviewed by Kaluzhny *et al.*, 2001). To determine whether the expression of other megakaryocyte-specific genes also depends on a physical interaction between GATA-1 and FOG-1, we analyzed megakaryocytes derived from *in vitro* differentiated murine embryonic stem (ES) cells and compared them with those derived from ES cells in which the GATA-1 gene had been replaced with the FOG-1-binding-defective GATA-1^{V205G} through homologous recombination (Chang *et al.*, 2002). ES cells were differentiated into embryoid bodies (EBs) in the presence of TPO to increase the number of megakaryocytes. AChE staining demonstrated that ~15% of the cells were megakaryocytes in both wild-type and mutant cultures (data not shown). Total RNA from these EBs was analyzed for α Ib gene expression by quantitative RT-PCR. The results show that α Ib levels in GATA-1^{V205G}-containing megakaryocytes were reduced to 16% when compared with their wild-type counterparts, consistent with a requirement for the GATA-1–FOG-1 interaction for α Ib gene expression *in vivo* (Figure 8). To determine whether other megakaryocyte-restricted genes similarly depend on the GATA-1–FOG-1 complex, we measured mRNA levels of the c-mpl, p45 NF-E2, GPIX, PF4 and PBP genes. Remarkably, GATA-1^{V205G}-containing megakaryocytes displayed a substantial reduction in the mRNA levels of p45 NF-E2, c-mpl and GPIX, but not of PF4 and PBP (Figure 8), indicating that the GATA-1–FOG-1 interaction is required for the expression of many, but not all, megakaryocyte-specific genes. Normal expression of the megakaryocyte-specific chemokines PF4 and PBP demonstrates that the GATA-1^{V205G}




Fig. 8. *In vivo* dependence of megakaryocyte-specific gene expression on physical interaction between GATA-1 and FOG-1. Total RNA was isolated from megakaryocyte-enriched EBs derived from *in vitro* differentiated wild-type or GATA-1^{V205G} mutant ES cells. mRNA levels of the indicated genes were determined using quantitative RT-PCR. Expression levels in GATA-1^{V205G} mutant cells were normalized for HPRT expression and plotted as a percentage of wild-type levels. The myeloid-expressed integrin gene α L served as additional control.

mutation did not simply abrogate megakaryocyte development. Instead, these results indicate that a subset of megakaryocyte-restricted genes is independent of a GATA-1–FOG-1 complex.

Activity of GATA-1 and FOG-1 on megakaryocytic gene promoters

Previous studies have shown that GATA-1 and FOG-1 synergistically activate the p45 NF-E2 promoter, similar to what we observed with the α IIb promoter (Tsang *et al.*, 1997). Therefore, we wanted to determine whether additional megakaryocytic gene promoters were also targets of GATA-1 and FOG-1. Specifically, we asked whether promoters of genes that are affected by the GATA-1^{V205G} mutation are activated by GATA-1 and FOG-1, and, conversely, whether genes that are expressed normally in GATA-1^{V205G}-containing megakaryocytes are independent of FOG-1. The c-mpl gene promoter has functionally important GATA- and Ets-binding elements within its first 100 bp of upstream sequence (Deveaux *et al.*, 1996). Therefore, we transfected a luciferase reporter gene construct driven by 100 bp of murine c-mpl promoter sequence into NIH-3T3 cells together with plasmids expressing GATA-1 and/or FOG-1. While GATA-1 alone activated the reporter gene 7-fold, co-expression of FOG-1 increased promoter activity up to 20-fold (Figure 9). In contrast, 222 bp of the murine PF4 promoter (Ravid *et al.*, 1991; Minami *et al.*, 1998), containing important GATA- and Ets-binding elements, showed activation by GATA-1, and but no further activation upon FOG-1 co-expression (Figure 9). Instead, FOG-1 repressed GATA-1 activity 6-fold, similar to what has been observed with the M1 α reporter gene. Preliminary results (not shown) further suggest that the PBP promoter is also not activated by GATA-1 and FOG-1. Together with the results obtained using the α IIb and p45 NF-E2 promoters, these data establish a correlation between gene activation by GATA-1 and




Fig. 9. GATA-1- and FOG-1-dependent activation of select megakaryocytic gene promoters. Transient transfection studies were performed as in Figure 1, with reporter gene constructs containing 100 bp of the murine c-mpl promoter or 222 bp of murine PF4 promoter. A promoterless (pGL2) vector served as negative control.

FOG-1 in transfection-based assays and sensitivity to the GATA-1^{V205G} mutation in megakaryocytes.

Discussion

GATA-1 and FOG-1 cooperate during the differentiation of megakaryocytes and erythroid cells (Tsang *et al.*, 1997; Crispino *et al.*, 1999; Nichols *et al.*, 2000; Chang *et al.*, 2002). FOG-1 binds to the N-terminal zinc finger of GATA-1, thereby modulating GATA-1 activity. The effects of FOG proteins on the transcriptional activities of GATA factors are determined by the cell type and the context of *cis*-regulatory elements present at a given promoter. To understand better the molecular basis for cooperative transcriptional activation by GATA-1 and FOG-1, we examined the effects of GATA-1 and FOG-1 on the α IIb gene promoter. We found that FOG-1 strongly increases GATA-1 activity on the α IIb promoter in a fashion that requires direct physical interaction. ChIP assays showed that GATA-1 and FOG-1 co-occupy this promoter in megakaryocytes *in vivo*, indicating that the actions of GATA-1 and FOG-1 are direct. Deletion analysis showed that 60 bp of the proximal promoter plus sequences containing the 5'-UTR were sufficient for full activation by GATA-1 and FOG-1 in NIH-3T3 cells. Remarkably, gain-of-function experiments, using synthetic promoter constructs, revealed that the mere presence of a specialized Ets element is sufficient to convert FOG-1 from a GATA-1 inhibitor into a co-activator. These findings provide a molecular explanation for the frequent occurrence of neighboring GATA and Ets elements in megakaryocyte-restricted genes.

Fli-1 probably functions at the proximal α IIb Ets element *in vivo* for several reasons. First, recombinant and cellular Fli-1 proteins bind to this site *in vitro* (Zhang *et al.*, 1993) (Figure 5B), consistent with the observation that the sequences flanking the proximal core Ets element match the Fli-1 consensus binding site perfectly (Zhang *et al.*, 1993; Mao *et al.*, 1994; Szymczyna and Arrowsmith, 2000). Secondly, ChIP experiments showed that Fli-1 binds the α IIb promoter *in vivo* in a megakaryocytic cell line (Figure 7). Thirdly, when fused

to GAL4, Fli-1 but not PU.1 and Spi-B can mediate transcriptional activation by GATA-1 and FOG-1. Fourthly, mice homozygous for a Fli-1 null mutation display altered megakaryocytic proliferation and differentiation (Hart *et al.*, 2000; Spyropoulos *et al.*, 2000; Kawada *et al.*, 2001). Finally, forced expression of Fli-1 in K562 cells augments α Ib expression (Athanasidou *et al.*, 1996), and Fli-1 can activate the α Ib promoter in transient transfection assays (Bastian *et al.*, 1999). However, it is likely that other Ets proteins might function at this site as well since anti-Fli-1 antibodies supershifted only part of the Ets protein complex that co-migrated with Fli-1 in primary megakaryocytes. Furthermore, transcriptional activation of the α Ib promoter by FOG-1 and GATA-1 was observed in NIH-3T3 cells, which do not express detectable amounts of Fli-1. Thus, select members of the large and diverse Ets protein family might substitute for Fli-1 function in non-hematopoietic tissues. Moreover, functional redundancy among Ets family proteins might explain the normal expression in Fli-1 null mice of several megakaryocytic genes that contain functional Ets elements in the regulatory regions, including c-mpl and α Ib (Hart *et al.*, 2000).

Through analysis of various Ets elements, we note that even when sequences do not match the Fli-1 consensus sequence, they might still be able to mediate activation by Fli-1. For example, based on sequence alone, the Ets sites in the GPIb α and GPIX promoters would be predicted to be poor binding sites for Fli-1. Yet both promoters are activated by Fli-1 and, in the case of GPIX, activation has been shown to depend on its Ets element (Bastian *et al.*, 1999). Therefore, promoter architecture appears to be critical for mediating transcriptional activation by Fli-1. In accordance with this interpretation, recruitment of Fli-1 but not PU.1 to the α Ib promoter via GAL4-binding sites led to activated transcription together with GATA-1 and FOG-1. These results suggest that megakaryocyte-specific promoter activity is determined by both the nucleotide sequence of the Ets-binding site and distinct domains within the bound Ets factor.

The above studies indicate that the proximal promoter region is sufficient for GATA-1- and FOG-1-dependent transcription, but they raise questions regarding the role of the distal conserved GATA and Ets sites between positions -457 and -506 bp upstream of the TSS. While both tandem GATA/Ets sites have enhancer activity, the distal enhancer is active in both erythroid and megakaryocytic lineages (Prandini *et al.*, 1992), indicating that it is not the critical determinant for megakaryocyte-specific expression of the α Ib gene. Furthermore, the distal Ets element does not conform to the Fli-1 consensus binding site. Indeed, PU.1, which has a substantially diverged DNA-binding domain and different binding site preference, has been shown to bind this Ets element (Doubekovski *et al.*, 1997). Megakaryocytes derived from PU.1-deficient ES cells expressed the α Ib gene normally, showing that PU.1 is not essential for α Ib expression (Zhang *et al.*, 1997). As appears to be the case for the proximal Ets site at -35 bp, it is possible that multiple Ets family members can bind and promote α Ib expression through the distal Ets element. The identity of these Ets proteins remains to be determined.

Direct physical interaction between GATA-1 and FOG-1 is required for α Ib expression in transfected cells and in ES cell-derived megakaryocytes, since a point mutation in GATA-1 that disrupts FOG-1 binding leads to loss of gene activation. Megakaryocytes bearing the GATA-1^{V205G} mutation express reduced levels of additional megakaryocyte-restricted genes, including c-mpl, p45 NF-E2 and GPIX, indicating that these genes are also regulated by the GATA-1-FOG-1 complex *in vivo*. In the cases of c-mpl and p45NF-E2, we and others (Tsang *et al.*, 1997) have shown that GATA-1 and FOG-1 can activate these genes in transfection-based assays, supporting a model in which GATA-1 and FOG-1 control these genes directly. It is important to note that the PF4 and PBP genes, which are expressed at late stages of megakaryocytic differentiation (Lepage *et al.*, 2000), were unaffected by the GATA-1^{V205G} mutation. This indicates that reduced expression of α Ib, c-mpl, p45 NF-E2 and GPIX is not simply the result of failed megakaryopoiesis, but instead reflects differences in GATA-1/FOG-1 dependence between distinct sets of genes. It is of note that the Ets elements in the upstream region of the c-mpl and GPIX genes have been shown to bind to Fli-1 *in vitro* (Deveaux *et al.*, 1996; Bastian *et al.*, 1999). Our observation that PF4 expression is independent of the GATA-1-FOG-1 interaction suggests alternative mechanisms of PF4 gene regulation. It is worth pointing out that the rat and human proximal PF4 promoters differ, with the human gene lacking a GATA-1 consensus element (Eisman *et al.*, 1990). Furthermore, the PF4 Ets elements do not conform to Fli-1 consensus sites, suggesting that other Ets proteins might regulate this gene. It is also possible that GATA-1 performs FOG-1-independent functions at this gene, or that GATA-2, which is expressed in megakaryocytes and also binds FOG-1 (Tsang *et al.*, 1997), might substitute for GATA-1. At the PBP gene, no functional GATA-1-binding sites have been identified, and GATA-1 did not activate the PBP promoter in transient transfection assays (M.Poncz, unpublished observation). Remarkably, all examined megakaryocyte-specific genes that were affected by the GATA-1^{V205G} mutation had regulatory regions that could be activated by GATA-1 and FOG-1 in transfection assays. In contrast, the promoters of the PF4 and PBP genes that were insensitive to the GATA-1^{V205G} mutation could not be activated by GATA-1 and FOG-1.

Our results are consistent with findings in patients with mutations in the N-terminal finger GATA-1 residues V205, G208 or D218 (Nichols *et al.*, 2000; Freson *et al.*, 2001; Mehaffey *et al.*, 2001). These patients exhibit different degrees of macrothrombocytopenia, bleeding diathesis and dyserythropoiesis. Among these GATA-1 mutations, V205M is the most severe. Patients carrying the V205M mutation were anemic and severely thrombocytopenic. Their bone marrows contained numerous dysplastic megakaryocytes, consistent with an essential role for the GATA-1-FOG-1 complex in megakaryopoiesis and platelet production. Since these patients had undergone bone marrow transplantation, the levels of various megakaryocyte-specific genes could not be measured.

It is instructive to compare our results with those obtained from megakaryocytes that express markedly reduced levels of wild-type GATA-1 as a result of a

targeted mutation in the GATA-1 regulatory region (Shivdasani *et al.*, 1997; Vyas *et al.*, 1999). While the GATA-1^{V205G} cells showed widespread, but not universal defects in megakaryocytic gene expression, the GATA-1-deficient megakaryocytes displayed reduced expression of all genes examined, including c-mpl, p45 NF-E2, α IIB and PF4. This more generalized decrease could be interpreted to reflect a general defect of megakaryocytic maturation. Alternatively, genes whose expression is diminished in the GATA-1-deficient but not in the GATA-1^{V205G} cells, might be regulated in a GATA-1-dependent, but FOG-1-independent, fashion. In addition, the degree to which GATA-2 might compensate for impaired GATA-1 activity might vary between GATA-1^{V205G}-containing and GATA-1-deficient megakaryocytes.

The frequent presence of GATA and Ets elements in megakaryocyte-expressed genes is indicative of the generality of our observations. This list of genes with functional GATA and Ets sites in their regulatory regions includes α IIB, c-mpl, GPIb α , GPV and GPIX, some of which are early markers of megakaryocytic development (Lepage *et al.*, 2000). Thus, we speculate that the GATA-1, FOG-1 and Fli-1 cooperativity is critical during the formation and/or maintenance of the megakaryocytic lineage. However, in the case of the p45 NF-E2 gene, it remains an open question whether GATA-1/FOG-1-mediated activation requires the presence of an Ets element, since transactivation experiments were performed on reporter gene constructs containing 7 kb of upstream region (Tsang *et al.*, 1997).

The mechanism by which Fli-1 exerts its effects on GATA-1 and FOG-1 remains to be determined. It is possible that Fli-1 forms specific contacts with FOG-1, thereby altering its conformation or promoting or interfering with additional protein contacts. It is also possible that Fli-1 communicates directly with GATA-1, although published studies showed relatively poor Fli-1–GATA-1 interaction *in vitro* (Rekhtman *et al.*, 1999). We found that amino acids 194–274 of Fli-1 were essential for its function in the GAL4 recruitment assays. This region contains a domain from amino acids 231 to 248 that is similar to the B-box present in select Ets proteins, including SAP1a and ELK1, but not in Ets-1, Ets-2 and PU.1 (Watson *et al.*, 1997). This domain has been shown to mediate contacts with other proteins to form higher order protein complexes on certain regulatory regions (Watson *et al.*, 1997). It is possible that other Ets proteins containing this domain might also be able to mediate GATA-1/FOG-1 synergy. This domain might aid in positioning the GATA-1–FOG-1 complex at the promoter in a way that is favorable for transcriptional activation. Alternatively, it might aid in replacing co-repressor complexes similar to what is observed in nuclear hormone receptors upon ligand binding (McKenna and O'Malley, 2002).

In summary, our work sheds new light on the mechanism by which megakaryocyte-specific gene expression is accomplished, providing new insights into the GATA–Ets signature motif found in the promoter regions of numerous megakaryocytic genes. In addition, we identified Fli-1 as a tissue-specific Ets-binding protein that converts FOG-1 from a repressor into a GATA-1 co-activator. The implications of these findings might extend to genes

controlled by GATA and FOG proteins in diverse tissues, including erythroid cells, lymphoid cells and the heart. Thus, we speculate that transcriptional regulators that cooperate with GATA factors in non-megakaryocytic cells might serve the same function as Fli-1 in megakaryocytes, which is to determine transcriptional output of GATA–FOG complexes.

Materials and methods

Plasmid constructs

The 912 bp rat α IIB promoter–growth hormone reporter construct has been described (Block *et al.*, 1994). Mouse α IIB promoter constructs were generated by PCR, and introduced between *Bgl*III and *Hind*III sites of the luciferase reporter vector pGL2-basic (Promega). The 946 bp mouse α IIB promoter was cloned into the human growth hormone-based vector pOGH (Nichols Institute Diagnostics) using the same restriction sites. The M1 α promoter (Martin and Orkin, 1990) was introduced into pGL2. Segments of 100 and 222 bp of the murine c-mpl and PF4 promoters, respectively, were subcloned into pGL2. In the pGL2– α IIB100 bp construct, the proximal Ets element between –32 and –50 bp was substituted with a single GAL4-binding site (5'-GCCGAGTAC-TGTCCTCCGA-3') by overlapping PCR (Block *et al.*, 1994). All PCR-based constructs were sequenced.

pXM-GATA-1, pXM-GATA-1^{V205G}, pMT2-FOG-1, pMFG-FOG-1^{S706R}, GAL4-PU.1^{1–160} and pCMX-GAL4 have been described (Martin and Orkin, 1990; Tsang *et al.*, 1997; Crispino *et al.*, 1999; Maitra and Atchison, 2000). FOG-1^{S706R} was subcloned into pMT2 at the *Eco*RI site. pCMX-GAL4-FLI-1^{1–274} and pCMX-GAL4-FLI-1^{1–194} were made by PCR and introduced into the GAL4 fusion expression vector pCMX.

Cell lines

NIH-3T3 cells were grown in Dulbecco's modified Eagle's medium (DMEM). Murine Y10 cells (Ishida *et al.*, 1993) were maintained in F-12 Nutrient Mixture (Invitrogen). All media contained 100 U/ml penicillin, 100 U/ml streptomycin, 2 mM L-glutamine and 10% fetal calf serum.

DNA transfection, reporter gene assay and western blotting

Cells were transfected with the calcium phosphate precipitation method. The total amount of transfected DNA was kept constant in all samples. After 48 h, luciferase activity was determined using commercial reagents (Promega). Growth hormone activity was assayed with a Nichols Diagnostic Institute kit.

In vitro ES cell differentiation, RNA extraction and semi-quantitative RT-PCR analysis

Wild-type and mutant TL-1 ES cell lines containing the GATA-1^{V205G} substitution have been described (Chang *et al.*, 2002). ES cells were differentiated into EBs as previously described using 100 ng/ml recombinant mouse TPO (R&D Systems) (Zhang *et al.*, 1997). Megakaryocytes in the EBs were identified by their distinct morphology and by their staining with AChE. Total cellular RNA was prepared using RNA STAT-60 reagent (TEL-TEST). The following sets of murine sense/antisense primers were used for RT-PCR: α IIB (GGCTGGAGCACACCTATGAGCT; GCTCAACCTGGGAGGCT); p45 NF-E2 (ACGTGGACATGTACCCAGTGG; GCCACCTGTCTTGCCCCGT); c-mpl (ACCAAGGTCCCTGGAGCG; AGGAGGCTGGGTCCACTT); GPIX (AGGCCCTGTACCTGCCAGTCC; GCCCAGCTCATAA-CCTGTGAGCT); PF4 (GTCCAGTGGCACCTCTGA; AATTGACATTTAGGCAGCTGA); PBP (GCCTGCCCACTTCATAACCTC; GGGTCCAGGCACGTTTTTTG); integrin α _L (GATCTGTACTACCTCATGGATCTC; GCAACTTGCAATTATGGCATCCAGC); and HPRT (TCCAGAAGTGGACACCTGC; GCTGGTGAAGAAGGACTCT). PCRs were performed in the presence of 1 μ Ci of [α -³²P]dCTP, separated on a polyacrylamide gel, and band intensities measured by Phosphor-imager analysis.

Chromatin immunoprecipitation assays

ChIP assays were performed as described (Forsberg *et al.*, 2000) using anti-GATA-1 (N6, Santa Cruz), mouse anti-Fli-1 (C-19, Santa Cruz), rabbit anti-Fli-1 (PharMingen) and affinity-purified rabbit anti-FOG-1 serum raised against amino acids 11–25 of mouse FOG-1. The following modifications were carried out: DNA–protein–antibody complexes were recovered by eluting twice with 100 μ l of 0.1 M NaHCO₃ and diluted with

TE buffer (10 mM Tris-HCl pH 7.4, 1 mM EDTA) to 400 μ l total volume. PCRs were performed for 26 cycles at 94°C for 20 s, 52°C for 30 s, 72°C for 45 s in the presence of [α - 32 P]dCTP using the following primer pairs: α Ib -110 bp to +226 bp region (GTCGACGTCTA-GAGGCTATTG; CTCTTAACCGCCCATATGTCCT); α Ib -3.3 to -3.6 kb region (TGTGAGTCCGCTGCCATT; TCTAGAGCAGGT-TAAGCCAG); and GAPDH (ACCACAGTCCATGCCATCAC; TCCACCACCTGTTGCTGTA). PCR products were separated on a 5% TBE polyacrylamide gel and quantitated by phosphorimager analysis.

Fetal liver cultures

Day 13 fetal livers from C57Blk6 mice were differentiated into morphologically mature megakaryocytes as described (Vyas *et al.*, 1999). AchE staining was performed after 5 days in culture with 100 ng/ml recombinant mouse TPO; cells were stained for AchE.

Electrophoretic mobility shift assays

A 0.2 ng aliquot ($\sim 10^5$ c.p.m.) of a [γ - 32 P]dATP- (Amersham) labeled probe was incubated with 10 μ g of nuclear extract. The sequence of the α Ib Ets probe was TAAGCTGAACTTCCGTTGGTGGGAAC, and that of the M1 α Ets probe was TAAGGATCAGCTTCCTCGAGC-GACCTT. A 2 μ g aliquot of the indicated antibodies: anti-FLI-1 (C19), anti-Ets-1 (N-276), anti-Ets-2 (C-20) and anti-PU.1 (T-21) (Santa Cruz Biotechnology) was added to nuclear extracts for 20 min prior to the addition of the probe.

Acknowledgements

We thank Stuart Orkin for the GATA-1^{V205G} ES cells, Mike Atchinson for the GAL4-PU.1 construct, and Merlin Crossley and Mitch Weiss for helpful discussions and critical comments on the manuscript. This work was supported by in part by grants PO1 HL40387 (M.P.) and RO1 DK54937 (G.A.B.), and gifts from the Schulman Foundation (M.P.) and the Plummer Family (M.P.).

References

Athanasidou, M., Clausen, P.A., Mavrothalassitis, G.J., Zhang, X.K., Watson, D.K. and Blair, D.G. (1996) Increased expression of the ETS-related transcription factor FLI-1/ERGB correlates with and can induce the megakaryocytic phenotype. *Cell Growth Differ.*, **7**, 1525–1534.

Bastian, L.S., Kwiatkowski, B.A., Breining, J., Danner, S. and Roth, G. (1999) Regulation of the megakaryocytic glycoprotein IX promoter by the oncogenic Ets transcription factor Fli-1. *Blood*, **93**, 2637–2644.

Block, K.L., Ravid, K., Phung, Q.H. and Poncz, M. (1994) Characterization of regulatory elements in the 5'-flanking region of the rat GPIIb gene by studies in a primary rat marrow culture system. *Blood*, **84**, 3385–3393.

Chang, A.N., Cantor, A.B., Fujiwara, Y., Lodish, M.B., Droho, S., Crispino, J.D. and Orkin, S.H. (2002) GATA-factor dependence of the multitype zinc-finger protein FOG-1 for its essential role in megakaryopoiesis. *Proc. Natl Acad. Sci. USA*, **99**, 9237–9242.

Crispino, J.D., Lodish, M.B., MacKay, J.P. and Orkin, S.H. (1999) Use of altered specificity mutants to probe a specific protein-protein interaction in differentiation: the GATA-1:FOG complex. *Mol. Cell*, **3**, 219–228.

Deveaux, S., Filipe, A., Lemarchandel, V., Ghysdael, J., Romeo, P.H. and Mignotte, V. (1996) Analysis of the thrombopoietin receptor (MPL) promoter implicates GATA and Ets proteins in the coregulation of megakaryocyte-specific genes. *Blood*, **87**, 4678–4685.

Doubeikovski, A., Uzan, G., Doubeikovski, Z., Prandini, M.-H., Porteu, F., Gisselbrecht, S. and Dusanter-Fourt, I. (1997) Thrombopoietin-induced expression of the glycoprotein IIb gene involves the transcription factor PU.1/Spi-1 in UT7-Mpl cells. *J. Biol. Chem.*, **272**, 24300–24307.

Eisman, R., Surrey, S., Ramachandran, B., Schwartz, E. and Poncz, M. (1990) Structural and functional comparison of the genes for human platelet factor 4 and PF4alt: characterization of the human β -thromboglobulin gene. Comparison with the gene for platelet factor 4. *Blood*, **76**, 336–344.

Forsberg, E.C., Downs, K.M., Christensen, H.M., Im, H., Nuzzi, P.A. and Bresnick, E.H. (2000) Developmentally dynamic histone acetylation pattern of a tissue-specific chromatin domain. *Proc. Natl Acad. Sci. USA*, **97**, 14494–14499.

Fossett, N. and Schulz, R.A. (2001) Conserved cardiogenic functions of the multitype zinc-finger proteins: U-shaped and FOG-2. *Trends Cardiovasc. Med.*, **11**, 185–190.

Fox, A.H., Liew, C., Holmes, M., Kowalski, K., Mackay, J. and Crossley, M. (1999) Transcriptional cofactors of the FOG family interact with GATA proteins by means of multiple zinc fingers. *EMBO J.*, **18**, 2812–2822.

Freson, K. *et al.* (2001) Platelet characteristics in patients with X-linked macrothrombocytopenia because of a novel GATA1 mutation. *Blood*, **98**, 85–92.

Fujiwara, Y., Browne, C.P., Cunniff, K., Goff, S.C. and Orkin, S.H. (1996) Arrested development of embryonic red cell precursors in mouse embryos lacking transcription factor GATA-1. *Proc. Natl Acad. Sci. USA*, **93**, 12355–12358.

Gaines, P., Geiger, J.N., Knudsen, G., Seshasayee, D. and Wojchowski, D.M. (2000) GATA-1 and FOG-dependent activation of megakaryocytic α Ib gene expression. *J. Biol. Chem.*, **275**, 34114–34121.

Graves, B.J. and Petersen, J.M. (1998) Specificity within the Ets family of transcription factors. *Adv. Cancer Res.*, **75**, 1–55.

Hart, A., Melet, F., Grossfeld, P., Chien, K., Jones, C., Tunnacliffe, A., Favier, R. and Bernstein, A. (2000) Fli-1 is required for murine vascular and megakaryocytic development and is hemizygotously deleted in patients with thrombocytopenia. *Immunity*, **13**, 167–177.

Ishida, Y., Levin, J., Baker, G., Stenberg, P.E., Yamada, Y., Sasaki, H. and Inoue, T. (1993) Biological and biochemical characteristics of murine megakaryoblastic cell line L8057. *Exp. Hematol.*, **21**, 289–298.

Kaluzhny, Y., Ravid, K. and Poncz, M. (2001) Transcription factors involved in lineage-specific gene expression during megakaryopoiesis. In Ravid, K. and Licht, J. (eds), *Transcription Factors: Normal and Malignant Development of Blood Cells*. Wiley-Liss, New York, NY, pp. 31–49.

Kawabata, H., Germain, R.S., Ikezoe, T., Tong, X., Green, E.M., Gombart, A.F. and Koeffler, H.P. (2001) Regulation of expression of murine transferrin receptor 2. *Blood*, **98**, 1949–1954.

Kawada, H., Ito, T., Pharr, P.N., Spyropoulos, D.D., Watson, D.K. and Ogawa, M. (2001) Defective megakaryopoiesis and abnormal erythroid development in Fli-1 gene-targeted mice. *Int. J. Hematol.*, **73**, 463–468.

Lemarchandel, V., Ghysdael, J., Mignotte, V., Rahuel, C. and Romeo, P.H. (1993) GATA and ETS cis-acting sequences mediate megakaryocyte-specific gene expression. *Mol. Cell. Biol.*, **13**, 668–676.

Lepage, A., Leboeuf, M., Cazenave, J.P., de la Salle, C., Lanza, F. and Uzan, G. (2000) The α (IIb) β (3) integrin and GPIb-V-IX complex identify distinct stages in the maturation of CD34(+) cord blood cells to megakaryocytes. *Blood*, **96**, 4169–4177.

Maitra, S. and Atchison, M. (2000) BSAP can repress enhancer activity by targeting PU.1 function. *Mol. Cell. Biol.*, **20**, 1911–1922.

Mao, X., Miesfeldt, S., Yang, H., Leiden, J.M. and Thompson, C.B. (1994) The FLI-1 and chimeric EWS-FLI-1 oncoproteins display similar DNA binding specificities. *J. Biol. Chem.*, **269**, 18216–18222.

Martin, D.I. and Orkin, S.H. (1990) Transcriptional activation and DNA binding by the erythroid factor GF-1/NF-E1/Eryf 1. *Genes Dev.*, **4**, 1886–1898.

McKenna, N.J. and O'Malley, B.W. (2002) Combinatorial control of gene expression by nuclear receptors and coregulators. *Cell*, **108**, 465–474.

Mehaffey, M.G., Newton, A.L., Gandhi, M.J., Crossley, M. and Drachman, J.G. (2001) X-linked thrombocytopenia caused by a novel mutation of GATA-1. *Blood*, **98**, 2681–2688.

Minami, T., Tachibana, K., Imanishi, T. and Doi, T. (1998) Both Ets-1 and GATA-1 are essential for positive regulation of platelet factor 4 gene expression. *Eur. J. Biochem.*, **258**, 879–889.

Nichols, K.E., Crispino, J.D., Poncz, M., White, J.G., Orkin, S.H., Maris, J.M. and Weiss, M.J. (2000) Familial dyserythropoietic anaemia and thrombocytopenia due to an inherited mutation in GATA1. *Nat. Genet.*, **24**, 266–270.

Pevny, L., Simon, M.C., Robertson, E., Klein, W.H., Tsai, S.-F., D'Agati, V., Orkin, S.H. and Costantini, F. (1991) Erythroid differentiation in chimaeric mice blocked by a targeted mutation in the gene for transcription factor GATA-1. *Nature*, **349**, 257–260.

Pevny, L., Lin, C.S., D'Agati, V., Simon, M.C., Orkin, S.H. and Costantini, F. (1995) Development of hematopoietic cells lacking transcription factor GATA-1. *Development*, **121**, 163–172.

Prandini, M.H., Uzan, G., Martin, F., Thevenon, D. and Marguerie, G. (1992) Characterization of a specific erythromegakaryocytic enhancer within the glycoprotein IIb promoter. *J. Biol. Chem.*, **267**, 10370–10374.

Ravid, K., Doi, T., Beeler, D.L., Kuter, D.J. and Rosenberg, R.D. (1991)

- Transcriptional regulation of the rat platelet factor 4 gene: interaction between an enhancer/silencer domain and the GATA site. *Mol. Cell. Biol.*, **11**, 6116–6127.
- Rekhtman,N., Radparvar,F., Evans,T. and Skoultchi,A.I. (1999) Direct interaction of hematopoietic transcription factors PU.1 and GATA-1: functional antagonism in erythroid cells. *Genes Dev.*, **13**, 1398–1411.
- Romeo,P.-H., Prandini,M.-H., Joulin,V., Mignotte,V., Prenant,M., Vainchenker,W., Marguerie,G. and Uzan,G. (1990) Megakaryocytic and erythrocytic lineages share specific transcription factors. *Nature*, **344**, 447–449.
- Shivdasani,R.A., Fujiwara,Y., McDevitt,M.A. and Orkin,S.H. (1997) A lineage-selective knockout establishes the critical role of transcription factor GATA-1 in megakaryocyte growth and platelet development. *EMBO J.*, **16**, 3965–3973.
- Spyropoulos,D.D., Pharr,P.N., Lavenburg,K.R., Jackers,P., Papas,T.S., Ogawa,M. and Watson,D.K. (2000) Hemorrhage, impaired hematopoiesis and lethality in mouse embryos carrying a targeted disruption of the *Fli1* transcription factor. *Mol. Cell. Biol.*, **20**, 5643–5652.
- Szymczyna,B.R. and Arrowsmith,C.H. (2000) DNA binding specificity studies of four ETS proteins support an indirect read-out mechanism of protein–DNA recognition. *J. Biol. Chem.*, **275**, 28363–28370.
- Tronik-LeRoux,D., Roullot,V., Schweitzer,A., Berthier,R. and Marguerie,G. (1995) Suppression of erythro-megakaryocytopoiesis and the induction of reversible thrombocytopenia in mice transgenic for the thymidine kinase gene targeted by the platelet glycoprotein α Ib promoter. *J. Exp. Med.*, **181**, 2141–2151.
- Tsang,A.P., Visvader,J.E., Turner,C.A., Fujiwara,Y., Yu,C., Weiss,M.J., Crossley,M. and Orkin,S.H. (1997) FOG, a multitype zinc finger protein, acts as a cofactor for transcription factor GATA-1 in erythroid and megakaryocytic differentiation. *Cell*, **90**, 109–119.
- Tsang,A.P., Fujiwara,Y., Hom,D.B. and Orkin,S.H. (1998) Failure of megakaryopoiesis and arrested erythropoiesis in mice lacking the GATA-1 transcriptional cofactor FOG. *Genes Dev.*, **12**, 1176–1188.
- Uzan,G., Prenant,M., Prandini,M., Martin,F. and Marguerie,G. (1991) Tissue-specific expression of the platelet GPIIb gene. *J. Biol. Chem.*, **266**, 8932–8939.
- Vyas,P., Ault,K., Jackson,C.W., Orkin,S.H. and Shivdasani,R.A. (1999) Consequences of GATA-1 deficiency in megakaryocytes and platelets. *Blood*, **93**, 2867–2875.
- Watson,D.K., Robinson,L., Hodge,D.R., Kola,I., Papas,T.S. and Seth,A. (1997) FLI1 and EWS–FLI1 function as ternary complex factors and ELK1 and SAP1a function as ternary and quaternary complex factors on the *Egr1* promoter serum response elements. *Oncogene*, **14**, 213–221.
- Weiss,M.J. and Orkin,S.H. (1995) GATA transcription factors: key regulators of hematopoiesis. *Exp. Hematol.*, **23**, 99–107.
- Zhang,C., Gadue,P., Scott,E., Atchison,M. and Poncz,M. (1997) Activation of the megakaryocyte-specific gene platelet basic protein (PBP) by the Ets family factor PU.1. *J. Biol. Chem.*, **272**, 26236–26246.
- Zhang,L., Lemarchandel,V., Romeo,P.H., Ben-David,Y., Greer,P. and Bernstein,A. (1993) The *Fli-1* proto-oncogene, involved in erythroleukemia and Ewing's sarcoma, encodes a transcriptional activator with DNA-binding specificities distinct from other Ets family members. *Oncogene*, **8**, 1621–1630.

Received May 2, 2002; revised July 25, 2002;
accepted August 14, 2002