

# The role of PACT in the RNA silencing pathway

Yoontae Lee<sup>1</sup>, Inha Hur<sup>1</sup>, Seong-Yeon Park<sup>1</sup>, Young-Kook Kim, Mi Ra Suh and V Narry Kim\*

Department of Biological Sciences and Research Center for Functional Cellulomics, Seoul National University, Seoul, Korea

**Small RNA-mediated gene silencing (RNA silencing) has emerged as a major regulatory pathway in eukaryotes. Identification of the key factors involved in this pathway has been a subject of rigorous investigation in recent years. In humans, small RNAs are generated by Dicer and assembled into the effector complex known as RNA-induced silencing complex (RISC) by multiple factors including hAgo2, the mRNA-targeting endonuclease, and TRBP (HIV-1 TAR RNA-binding protein), a dsRNA-binding protein that interacts with both Dicer and hAgo2. Here we describe an additional dsRNA-binding protein known as PACT, which is significant in RNA silencing. PACT is associated with an ~500 kDa complex that contains Dicer, hAgo2, and TRBP. The interaction with Dicer involves the third dsRNA-binding domain (dsRBD) of PACT and the N-terminal region of Dicer containing the helicase motif. Like TRBP, PACT is not required for the pre-microRNA (miRNA) cleavage reaction step. However, the depletion of PACT strongly affects the accumulation of mature miRNA *in vivo* and moderately reduces the efficiency of small interfering RNA-induced RNA interference. Our study indicates that, unlike other RNase III type proteins, human Dicer may employ two different dsRBD-containing proteins that facilitate RISC assembly.**

*The EMBO Journal* (2006) 25, 522–532. doi:10.1038/sj.emboj.7600942; Published online 19 January 2006

Subject Categories: RNA

Keywords: Dicer; microRNA; PACT; RNAi; siRNA

## Introduction

Small RNAs function as guide molecules in a wide range of regulatory pathways that are now collectively known as RNA silencing (Bartel, 2004; Baulcombe, 2005). Through specific base-pairing with mRNAs, these tiny ~22 nt RNAs induce mRNA degradation and translational repression. Based on their origin, small RNAs can be grouped into two classes: microRNA (miRNA) or small interfering RNA (siRNA) (Kim, 2005b). miRNAs originate from endogenous transcripts harboring local hairpin structures, while siRNAs are generated from long dsRNAs. siRNAs can be further classified into

transacting siRNA (tasiRNA), repeat-associated siRNA (rasiRNA), and small scan RNA (scnRNA).

miRNAs are transcribed from endogenous genes by RNA polymerase II (Cai *et al.*, 2004; Lee *et al.*, 2004a; Kim, 2005a). The hairpin embedded in the primary transcripts of miRNA (pri-miRNAs) are cropped into hairpin-structured precursors (pre-miRNAs) by the nuclear RNase III Drosha (Lee *et al.*, 2003). Pre-miRNAs are then exported to the cytoplasm with the help of exportin-5 (Exp5) (Yi *et al.*, 2003; Bohnsack *et al.*, 2004; Lund *et al.*, 2004) and subsequently processed into mature miRNA of ~22 nt by another RNase III Dicer (Grishok *et al.*, 2001; Hutvagner *et al.*, 2001). Unlike miRNAs, siRNAs are generated from long double-stranded RNAs (dsRNAs) by Dicer (Bernstein *et al.*, 2001; Ketting *et al.*, 2001). Thus, Dicer serves as a common enzyme in the biogenesis of both miRNAs and siRNAs in mammalian cells.

The small RNAs are loaded onto the effector complex, known as the RNA-induced silencing complex (RISC) (Filipowicz, 2005; Sontheimer, 2005; Tomari and Zamore, 2005; Zamore and Haley, 2005). Argonaute proteins are the core components of the RISC (Hammond *et al.*, 2001; Carmell *et al.*, 2002; Liu *et al.*, 2004; Meister *et al.*, 2004; Sontheimer and Carthew, 2004). Argonaute proteins have three highly conserved regions: the PAZ domain at the N-terminus, the middle conserved domain, and the PIWI domain at the C-terminus. Structural and biochemical studies indicate that Argonaute proteins are capable of interacting directly with small RNAs through their PAZ domains (to the 3' end) and through the PIWI and the middle domains (to the 5' end) (Song *et al.*, 2003, 2004; Yan *et al.*, 2003; Lingel *et al.*, 2004; Ma *et al.*, 2004, 2005; Parker *et al.*, 2005; Rivas *et al.*, 2005; Yuan *et al.*, 2005). A small RNA guides the RISC to its target RNA, which leads to mRNA degradation and/or to translational repression.

The mechanism of RISC formation has been intensively studied in several types of organisms, such as flies and humans (Filipowicz, 2005; Sontheimer, 2005; Tomari and Zamore, 2005; Zamore and Haley, 2005). In *D. melanogaster*, there are two types of Dicers, Dicer-1 and Dicer-2 (Lee *et al.*, 2004b). Dicer-1 is responsible for miRNA biogenesis and possibly for the assembly of miRNA-incorporated RISC (miRISC), while Dicer-2 is able to process only siRNAs from long dsRNAs. Dicer-2 also appears to be required for loading a guide strand of siRNA duplex onto the RISC complex (siRISC). There is only one type of Dicer in humans, which generates both miRNA and siRNA. Besides the processing of miRNA, human Dicer has also been proposed to have a role in RISC assembly, because a depletion of Dicer results in a defect of RNA interference (RNAi) process mediated by siRNA duplex in human cell lines (Doi *et al.*, 2003).

Several dsRNA-binding proteins are known to associate with RNase III proteins to promote miRNA processing or RISC programming. In the nucleus, Drosha associates with DGCR8 (in humans) or Pasha (in *Drosophila* and *Caenorhabditis elegans*), which contains tandem dsRNA-binding domains (dsRBDs), to form the microprocessor complex that processes

\*Corresponding author. Institute of Biological Sciences and Research Center for Functional Cellulomics, Seoul National University, Seoul 151-742, Korea. Tel.: +82 2 880 9120; Fax: +82 2 887 0244; E-mail: narrykim@snu.ac.kr

<sup>1</sup>These authors contributed equally to this work

Received: 8 November 2005; accepted: 9 December 2005; published online: 19 January 2006

pri-miRNAs into pre-miRNAs (Denli *et al*, 2004; Gregory *et al*, 2004; Han *et al*, 2004; Landthaler *et al*, 2004). Loquacious (Loqs) (also known as R3D1), which has three dsRBDs, is a partner of *Drosophila* Dicer-1 in flies (Forstemann *et al*, 2005; Jiang *et al*, 2005; Saito *et al*, 2005). The Dicer-1–Loqs heterodimer efficiently carries out pre-miRNA processing. When Loqs was depleted by RNAi in the *Drosophila* S2 cell line, the mature miRNA level was reduced, while pre-miRNAs accumulated, indicating that Loqs is required for the pre-miRNA cleavage event (Forstemann *et al*, 2005; Jiang *et al*, 2005; Saito *et al*, 2005). R2D2, another dsRNA-binding protein, is known to associate with *Drosophila* Dicer-2 to promote siRNA from entering into Ago2 (Liu *et al*, 2003). Dicer-2 and R2D2 were found in holo-RISC and in the RISC-loading complex (RLC) of the *Drosophila* cell extract (Pham *et al*, 2004; Tomari *et al*, 2004a). The heterodimer of Dicer-2 and R2D2 is capable of selecting a guide strand from siRNA duplex by sensing the thermodynamic stability of both ends of the siRNA duplex (Tomari *et al*, 2004b). Recently, TRBP (HIV-1 TAR RNA-binding protein), a human homolog of Loqs, was reported to interact with Dicer and human Ago2 (hAgo2). The depletion of TRBP by RNAi causes defects of siRNA- or miRNA-mediated RNA silencing processes in human cell lines (Chendrimada *et al*, 2005; Haase *et al*, 2005).

PACT is similar to TRBP in the domain structure, implicating that PACT may be another homolog of Loqs (Saito *et al*,


2005). PACT was initially known as a protein activator of PKR (Patel and Sen, 1998). Through its dsRBD1 and dsRBD2, PACT interacts with PKR (Peters *et al*, 2001). The dsRBD3 at the C-terminus of PACT functions as a PKR activation domain (Peters *et al*, 2001). Contrary to PACT, TRBP was proposed as an inhibitor of PKR. TRBP interacts with PKR in a very similar manner with PACT, except that its dsRBD3 functions as the PKR inhibition domain, instead of as the activation domain (Gupta *et al*, 2003). Thus, PACT and TRBP have opposite roles in the regulation of PKR, although their domain structures are very similar to each other.

In this study, we explore the role of dsRNA-binding proteins in RNA silencing and identify PACT as a novel component of human RISC.

## Results

### PACT interacts with Dicer

To identify the cofactor for Dicer in humans, various dsRNA-binding proteins were fused to V5 tag and coexpressed with FLAG-Dicer protein in HEK293T cells for coimmunoprecipitation and Western blot analysis. Among these proteins, we found that PACT and TRBP interacted efficiently with Dicer (Figure 1A), whereas DGCR8, SPNR, Staufen, and two hypothetical dsRNA-binding proteins (FLJ20036 and FLJ20399) did not bind significantly to Dicer (data not shown). Immunocytochemistry for PACT, TRBP, and Dicer fused to V5 tag



**Figure 1** PACT interacts with Dicer, but does not facilitate the cleavage activity. (A) Immunoprecipitation followed by Western blotting. V5-Dicer protein was coexpressed in HEK293T cells with FLAG-PACT, or FLAG-TRBP protein. Immunoprecipitation was carried out using anti-FLAG antibody and the protein was visualized using either anti-V5 or anti-FLAG antibody. (B) *In vitro* pre-miRNA processing assay. Total HEK293 T extract (total lysate) or immunoprecipitates (FLAG-IP) of FLAG-Dicer, FLAG-TRBP, or FLAG-PACT were incubated with 5' end radiolabeled pre-let-7a-1. The RNA was extracted by phenol extraction method, followed by separating on 15% urea-PAGE. (C) *In vitro* processing assay using recombinant human Dicer and PACT. In all, 200 ng of recombinant PACT prepared from *E. coli* and 0.25 U of recombinant human Dicer (Stratagene) were used in each reaction. Pre-let-7a-1 (left panel) and long dsRNA of 80 bp (right panel) were used as substrates.



the recent study on TRBP using yeast two hybrid assay by Haase *et al* (2005).

Next, we looked for the region in Dicer that is required for binding to PACT or TRBP. Four deletion mutants were generated ( $\Delta$ DEAD,  $\Delta$ HelicC,  $\Delta$ DUF, and  $\Delta$ PAZ) (Figure 2A). The Dicer mutants,  $\Delta$ HelicC,  $\Delta$ DUF, and  $\Delta$ PAZ, were unable to interact with either PACT or TRBP. Thus, the highly conserved helicase motif whose biochemical role in Dicer remains unknown may be critical for the interaction with PACT and TRBP (Figure 2C and D).

Interestingly, when the Dicer mutant proteins were immunoprecipitated and assayed for pre-miRNA processing activity,  $\Delta$ HelicC mutant lacking the helicase motif was capable of processing (Figure 2E, lane 4). This result shows that the helicase motif is dispensable for processing and that PACT and TRBP may not be required for the Dicer cleavage reaction itself. It is noted that the DUF283 domain whose function remains unknown may be critical for processing, as the mutant lacking this domain ( $\Delta$ DUF) lost the activity (Figure 2E, lane 5).


### PACT binds to hAgo2

Both Dicer and TRBP have previously been shown to interact with hAgo2 (Chendrimada *et al*, 2005). To test whether PACT also associates with hAgo2, FLAG-PACT protein was coexpressed with V5-hAgo2 (Figure 3). Coimmunoprecipitation followed by Western blotting revealed that PACT is indeed associated with hAgo2 (Figure 3A). The dsRBD3 mutant of PACT is capable of binding to hAgo2, indicating that PACT may bind to hAgo2 independently of Dicer (Figure 3B). Like PACT, the mDR3 mutant of TRBP bound to hAgo2 efficiently (data not shown), suggesting that the mode of interaction between TRBP and hAgo2 is similar to that between PACT and hAgo2.

### PACT, TRBP, and Dicer are associated with the slicer activity in vitro


We presented in Figure 1B that PACT and TRBP are associated with pre-miRNA processing activity, indicating that the complex(es) with Dicer may be preassembled prior to the pre-miRNA processing step. A similar *in vitro* processing experiment was performed by incubating immunoprecipitated FLAG-hAgo2 protein with 5' end-labeled pre-miR-30a (Figure 4A, lane 2). Intriguingly, the FLAG-hAgo2 immunoprecipitate was also capable of executing pre-miRNA processing. This result supports the recent findings that hAgo2 may be preassembled in the pre-miRNA processing complex in human cells (Gregory *et al*, 2005; Maniataki and Mourelatos, 2005; Meister *et al*, 2005). Further supporting this notion, when the beads were washed after processing reaction, the miRNA product was found to be associated with hAgo2, Dicer, TRBP, and PACT (Figure 4A, lanes 6–10).

Next, we asked if PACT and TRBP are present in the active RISC using the *in vitro* RNAi protocol that was recently developed by Maniataki and Mourelatos (2005) (Figure 4B). The FLAG-fused proteins were expressed in HEK293T cells and immobilized on anti-FLAG beads. The immunoprecipitates were then incubated with cold pre-miR-30a to allow pre-miRNA processing. As a control, 27-mer siRNA, which does not contain a complementary sequence to target RNA, was used. The beads were washed twice to remove excess pre-miRNA and then incubated with labeled target RNA, which is complementary to miR-30a-5p. Not only FLAG-hAgo2, but also FLAG-Dicer, FLAG-TRBP and FLAG-PACT mediated target cleavage reaction (Figure 4B). This result implies that PACT and TRBP are associated with hAgo2 in the functional RISC. Therefore, hAgo2, Dicer, PACT and TRBP may be preassembled prior to pre-miRNA processing and remain together to the final RISC, although the stoichiometry and the conformation of the complex are likely to change over



**Figure 3** PACT binds to hAgo2. (A) Immunoprecipitation followed by Western blotting. FLAG-PACT or FLAG-TRBP proteins, V5-Dicer and V5-hAgo2 protein were coexpressed in HEK293T cells. Immunoprecipitation was carried out using anti-FLAG antibody and the protein was visualized using either anti-V5 or anti-FLAG antibody. (B) The same experiment as described in (A), except for using FLAG-PACT mutants and V5-hAgo2 (wild type). In this experiment, V5-Dicer was not coexpressed.





**Figure 5** PACT and TRBP coexist in the ~500 kDa RISC. Total extract was prepared from HEK293T cells transiently transfected with FLAG-PACT, V5-Dicer, and V5-hAgo2 expression vectors and fractionated through a Sephacryl-S300 HR column. Each fraction was subject to immunoprecipitation with anti-FLAG antibody. The immunoprecipitate was used for (A) Western blot analysis using anti-TRBP or anti-FLAG antibody, or (B) *in vitro* RNAi assay using miR-30a-5p target RNA. The fraction number and the protein molecular mass standard (Sigma) are indicated at the top of the figures.

capable of pre-miRNA processing and target cleavage. This complex may be comparable to the Dicer-TRBP-hAgo2 complex described recently by Shiekhhattar and colleagues (Chendrimada *et al*, 2005; Gregory *et al*, 2005).

#### **PACT is required for siRNA-induced RNAi**


To examine whether PACT is required for RNAi reaction induced by siRNA, we transfected siRNA against PACT, TRBP, and Dicer together with siRNA against luciferase and the luciferase reporter constructs (Figure 6A). siRNA against green fluorescence protein (siGFP) was used as a control. Depletion of PACT resulted in the reduction of the efficiency of RNAi against luciferase (Figure 6A). Although the effect of RNAi against TRBP was reproducibly stronger than that against PACT (Figure 6A), our data indicate that PACT may also be required for RNA silencing pathway.

RNAi against TRBP did not significantly affect the level of Dicer protein, which is consistent with the finding by Haase *et al* (2005) (Figure 6B). When PACT was depleted, the level of Dicer decreased slightly, suggesting that PACT, but not

TRBP, may contribute to the stabilization of Dicer protein (Figure 6B).

#### **PACT is required for the accumulation of mature miRNA**

We then investigated whether PACT has any responsibility for miRNA biogenesis *in vivo* (Figure 7A). PACT, TRBP, Dicer, or exportin-5 was depleted by RNAi from a HeLa cell line that expresses pri-miR-30a from the tetracycline-inducible promoter. Reduction of the target mRNA (PACT, TRBP, Dicer, exportin-5) was verified by RT-PCR (Figure 7C). After incubation with siRNA, the cell line was exposed to doxycycline, the derivative of tetracycline, for induction of pri-miR-30a. Northern blotting on miR-30a-5p indicated that PACT may be required for the accumulation of newly synthesized miRNA. Interestingly, the depletion of PACT resulted in stronger effects on miRNA compared to that of TRBP. We also generated an additional HeLa cell line that expresses pri-miR-124a from the tetracycline-inducible promoter (Figure 7B). Accumulation of miR-124a was dependent on PACT in this cell line.



**Figure 6** PACT is required for siRNA-induced RNAi. (A) HeLa cells were transiently transfected with siRNA targeting Dicer, PACT, or TRBP along with siRNA against firefly luciferase, firefly luciferase-expression plasmid, and renilla luciferase-expression plasmid. The firefly luciferase activity was normalized against renilla luciferase activity. This experiment was performed in triplicate. Error bars indicate standard deviations. (B) Western blot analysis. siRNA targeting GFP, Dicer, hAgo2, TRBP, or PACT was transfected into HeLa cells. Following transfection (48 h), total extract was prepared in lysis buffer (10 mM Tris, pH 7.4, 1 mM EDTA, 500 mM NaCl, 0.5% Triton-X100). In all, 120  $\mu$ g of total extract was loaded on 10% SDS-PAGE. Endogenous Dicer, TRBP, or PACT was visualized using anti-Dicer, anti-TRBP, or anti-PACT antibody.

It is noted that the endogenous levels of miR-16 and let-7a in unmodified HeLa cells were changed only slightly after the depletion of PACT (data not shown). When TRBP was knocked down in the same condition, we did not observe significant changes in miRNA level (data not shown), which is consistent with the data of Filipowicz and colleagues (Haase *et al*, 2005). This also supports the previous observations that mature miRNA is highly stable in cells and therefore the changes in the steady-state level of mature miRNA may be difficult to detect (Lee *et al*, 2003; Yi *et al*, 2003; Lund *et al*, 2004).

## Discussion

In this study, we present PACT as another dsRNA-binding protein that functions as a component of the human RISC.


Human Dicer may associate with two different dsRNA-binding proteins, PACT and TRBP. Our *in vitro* pre-miRNA processing assay using a Dicer mutant lacking the helicase motif indicates that Dicer does not need a cofactor for pre-miRNA cleavage (Figure 2). However, the depletion of PACT by RNAi resulted in a reduction of mature miRNAs (Figure 7). These seemingly paradoxical data can be explained if PACT has a function in RISC formation. If a step of RNA loading into Argonaute protein is blocked, these small RNAs may become unstable. In fact, when PACT was depleted in HeLa cells, the decrease of mature miRNA, but not the accumulation of pre-miRNA, was monitored by Northern blot analysis (Figure 7). Our results suggest that PACT may play a role at a certain step of the RISC assembly, but not at the pre-miRNA cleavage step.

In flies, two different dsRBD proteins, Loqs/R3D1 and R2D2, interact with Dicer-1 and Dicer-2, respectively. Loqs/R3D1 facilitates pre-miRNA cleavage by Dicer-1, while R2D2 is not required for dsRNA cleavage, but instead participates in strand selection. The deletion of PRBP, a mouse homolog of TRBP, yields viable mice that often die at weaning, while knocking out Dicer results in embryonic lethality (Zhong *et al*, 1999; Bernstein *et al*, 2003), implicating that in mice, PRKRA, a mouse homolog of PACT, may compensate for PRBP during development (Chendrimada *et al*, 2005; Haase *et al*, 2005).

Our study supports this possibility that, in humans, the roles of dsRBD-containing Dicer cofactors may be partially redundant because RNAi experiments indicate that both PACT and TRBP may be involved in siRNA-induced RNAi and in miRNA accumulation. They also behave similarly in the modes of protein interaction: (1) two proteins share the common binding region in Dicer (246–584 a.a.), and (2) the third dsRBDs of PACT and TRBP are important for binding to Dicer.

However, our data also suggest that the roles of PACT and TRBP may be differentiated to some extent. The depletion of PACT resulted in a strong effect on miRNA accumulation (Figure 7) and a relatively moderate effect on RNAi reporter assay (Figure 6). In contrast, RNAi against TRBP had only limited effects on miRNA level, whereas the same siRNA induced clear effects on RNAi efficiency. One plausible possibility is that PACT is more suitable for the role in the transition step of miRNA duplex into single-strand miRNA, while TRBP is more active in the later stages of RISC formation and target cleavage. Thus, the roles of PACT and TRBP may be differentiated in normal conditions, but they may act redundantly to some extent in the absence of the other.

Our results from the coupled *in vitro* processing-RNAi assay support and extend the findings by Maniataki and Mourelatos (2005) that the protein components of the RISC complex seem to be preassembled prior to pre-miRNA processing in humans (Figure 4). Comparable results were reported recently by other groups (Gregory *et al*, 2005; Maniataki and Mourelatos, 2005; Meister *et al*, 2005). Gel filtration and coimmunoprecipitation experiments revealed that hAgo2, Dicer, TRBP, and PACT interact with one another, forming the RISC of ~500 kDa (Figure 5). In *Drosophila*, the assembly of the RISC appears to be achieved in a stepwise manner. The Dicer-2–R2D2 heterodimer binds to siRNA duplex to form the R1 complex that is later converted into the intermediate complex R2/RLC (RISC loading complex). Subsequently, R2/RLC recruits Ago2 to form the mature



**Figure 7** PACT is required for the accumulation of mature miRNA. (A) Northern blot analysis. siRNA targeting Dicer, TRBP, or PACT was transfected into a HeLa cell line that expresses primary miR-30a under the control of the tetracycline-inducible promoter. Following transfection (48 h), doxycycline (3 µg/ml) was added to the medium to induce the expression of miR-30a. After 24 h, the cells were harvested for Northern blotting (the left panel). After normalization for loading, the relative ratios for mature miR-30a were calculated and normalized to no doxycycline-treated experiment (lane 1). The average value obtained from two independent experiments was presented in the right panel. (B) The same experiment as described in (A), except for using pri-miR-124a inducible HeLa cell line. The left panel shows a representative Northern blot result. The right panel presents the mean value of relative expression level for mature miR-124a obtained from two independent experiments. (C) RT-PCR. Total RNA used in (A) was reverse-transcribed and amplified by PCR using gene-specific primers.

RISC, which likely acts as the RNAi effector in *Drosophila* (Pham and Sontheimer, 2005). Our data indicate that the mechanism of human RISC formation may be different from that of flies. It is also possible that the siRISC assembly process may differ from that of miRISC.

PACT and TRBP are known as regulators of PKR. PKR is a dsRNA-dependent serine/threonine protein kinase, which phosphorylates eIF2 $\alpha$  on Ser51 to cause a general reduction of protein synthesis (Taylor *et al*, 2005). Besides eIF2 $\alpha$ , several proteins are known to be phosphorylated by PKR,

such as NFAR-1, NFAR-2, and human protein phosphatase 2A (PP2A) regulatory subunit B56 $\alpha$  (Xu and Williams, 2000; Saunders *et al*, 2001). It is possible that the component(s) of RISC may be regulated by PKR through phosphorylation and that PACT and TRBP may regulate PKR activity to control the RISC activity. Alternatively, a competition between the RNA silencing pathway and the PKR pathway may exist, because the third dsRBDs of both PACT and TRBP are responsible for their interaction with Dicer as well as for the regulation of PKR. It would be intriguing to investigate

the possibility of the crosstalk between the RNA silencing pathway and the PKR pathway.

## Materials and methods

### Cloning of PACT and TRBP

PCR products of PACT and TRBP were subcloned into FLAG-pCK vector for expression in human cells, at the *Bam*HI and *Xho*I sites. Primer sequences used for PCR are as follows. For PACT, 5'-GGATCCATGTCAGAGCAGGCACC-3' (forward) and 5'-CTCGAGCTCCAGATTTACTTTCTTCTGTC-3' (reverse) were used. For TRBP, 5'-GGATCCATGAGTGAAGAGGAGCAAGGCTC-3' (forward) and 5'-CTCGAGCTCCAGATTTACTTTCTTCTGTC-3' (reverse) were used.

### Mutagenesis of PACT and Dicer

Point mutagenesis of PACT was carried out by using a site-directed mutagenesis kit (Stratagene) according to the manufacturer's manual. Primer sequences used for mutagenesis are as follows. For PACT mDR1 mutant, 5'-CTGGCGAAACATAGAAAGAAGGAGGCTGCCATAAAC-3' (upper primer) and 5'-GTTTATGGCAGCCTCCTTCTTCTATGTTTCGCCAG-3' (lower primer) were used. For PACT mDR2 mutant, 5'-GCAAGCCAAAAGGAATAAGAAGGAGAAATTTCTTGCC-3' (upper primer) and 5'-GGCAAGAAATTTCTCCTTCTTATCCTTTGGCTTGC-3' (lower primer) were used. For PACT mDR3 mutant, 5'-GCAATGCACAAAGTGATAAGAAGCACAATGCTTTGCCAG-3' (upper primer) and 5'-CTGCAAAGCATTGTGCTTCTTATCACTTTGTGCATTCG-3' (lower primer) were used.

To prepare Dicer deletion mutants, PCR products of Dicer deletion mutants were subcloned into FLAG-pcDNA3.1 vector (Invitrogen) for expression in human cells, at the *Hind*III and *Apa*I sites. Primer sequences used for PCR are as follows. For  $\Delta$ DEAD, 5'-AAGCTTTATACTTCTCAGCCATGTGAG-3' (forward) and 5'-CACAGTCGAGGCTGATCAG-3' (reverse) were used. For  $\Delta$ HelicC, 5'-AAGCTTATCTTGAGAAACAAGTGTCC-3' (forward) and 5'-CACAGTCGAGGCTGATCAG-3' (reverse) were used. For  $\Delta$ DUF, 5'-AAGCTTATTCCAGAGTGTGTTGAGGGATAG-3' (forward) and 5'-CACAGTCGAGGCTGATCAG-3' (reverse) were used. For  $\Delta$ PAZ, 5'-AAGCTTTACCTAACTTAGACTTCG-3' (forward) and 5'-CACAGTCGAGGCTGATCAG-3' (reverse) were used.

### Cell culture and transfection

HeLa cells and HEK293T cells were cultured in DMEM (WelGENE) supplemented with 10% FBS (WelGENE). Transfection was carried out by calcium-phosphate method.

### Immunoprecipitation

HEK293T cells grown in a 10-cm dish were collected in 500  $\mu$ l of ice-cold buffer D-K'100 (20 mM Tris, pH 8.0, 100 mM KCl, 0.2 mM EDTA, 0.2 mM PMSF) 48 h post-transfection. The cells were sonicated on ice and centrifuged at 13 200 r.p.m. for 15 min at 4°C. The supernatant was incubated with 10  $\mu$ l of anti-FLAG antibody conjugated to agarose beads (anti-FLAG M2 affinity gel, Sigma) with constant rotation for 60 min at 4°C. The beads were washed five times in buffer D-K'100, drained, and used for Western blot analysis.

### In vitro RNAi assay

HEK293T cells grown in two 10-cm dishes were collected in 500  $\mu$ l of ice-cold buffer D-K'100 48 h post-transfection. The cells were sonicated on ice and centrifuged at 13 200 r.p.m. for 15 min at 4°C. The supernatant was incubated with 20  $\mu$ l of anti-FLAG antibody conjugated to agarose beads (anti-FLAG M2 affinity gel, Sigma) with constant rotation for 2 h at 4°C. The beads were washed three times in buffer D-K'100, and were washed three times in RNAi reaction buffer (30 mM HEPES, pH 7.5, 40 mM potassium acetate, 5 mM magnesium acetate, 5 mM DTT). The processing reactions were performed in a total volume of 20  $\mu$ l that consisted of 15  $\mu$ l of the beads from immunoprecipitation, 1  $\mu$ l of ribonuclease inhibitor (40 U/ $\mu$ l, TAKARA), 3  $\mu$ l of RNAi reaction buffer, and 1  $\mu$ l of synthetic pre-miR-30a (10  $\mu$ M). As a control, 1  $\mu$ l of 27mer siRNA (10  $\mu$ M) was used. The sequences of 27mer siRNA are 5'-CCAUGGUAUCCGAAAGUUAGGCUdGdG-3' (sense) and 5'-CCAGCCUAACUUUCCGUAUCAUGGUA-3' (antisense). The reaction mixture was incubated at 37°C for 90 min. Before target RNA cleavage reaction, the beads were washed twice with RNAi reaction buffer. In all, 1  $\mu$ l

of ribonuclease inhibitor (TAKARA), 0.5  $\mu$ l of the radiolabeled target RNA (miR-30a-5p target RNA,  $1 \times 10^4$ – $1 \times 10^5$  c.p.m.), and 0.1  $\mu$ l of yeast RNA (1  $\mu$ g/ $\mu$ l, Ambion) were added to 20  $\mu$ l of the washed beads. The reaction mixture was incubated at 37°C for 90 min. RNA was extracted from the reaction mixture by phenol extraction and analyzed on 15% denaturing urea polyacrylamide gel. The sequence of miR-30a-5p target RNA is 5'-AACCUUGCUUCCAGUCGAGGAUGUUUACACCAAG-3'. Pre-miR-30a, 27mer control siRNA and miR-30a-5p target RNA were purchased from Samchully Pham Co. Ltd. The miR-30a-5p target RNA was labeled at the 5' end using T4 polynucleotide kinase (TAKARA) and [ $\gamma$ - $^{32}$ P]ATP.

### Gel exclusion chromatography

Gel exclusion chromatography was carried out as described previously (Han *et al*, 2004), with the following modifications. HEK293T cells grown in 20 10-cm dishes were cotransfected with FLAG-PACT, V5-Dicer, and V5-hAgo2 expression vectors. Total cell extract was prepared in buffer D'-K100 by sonication and centrifugation, and then concentrated to the final volume of 1 ml using Centricon YM-30 (Millipore). This total extract was treated with 10  $\mu$ l of 10 mg/ml RNase A and incubated at 4°C for 30 min. This extract was loaded on a Sephacryl S-300 HR (Sigma) column and fractionated. Each fraction was 1.65 ml in volume. In all, 1.6 ml of each fraction was taken for immunoprecipitation with anti-FLAG antibody, and the immunoprecipitate was used for Western blot analysis using anti-TRBP or anti FLAG antibody or for *in vitro* RNAi assay.

### In vitro processing of pre-miRNAs

*In vitro* processing of pre-miRNAs using FLAG-immunoprecipitates or recombinant human Dicer (Stratagene) was carried out as described previously (Lee *et al*, 2002, 2003). Briefly, 30  $\mu$ l of processing reaction contained 3  $\mu$ l of 64 mM MgCl<sub>2</sub>, 1 U/ $\mu$ l of ribonuclease inhibitor (TAKARA), the radiolabeled pre-miRNA of  $1 \times 10^4$ – $1 \times 10^5$  c.p.m., and 15  $\mu$ l of the beads from immunoprecipitation. In Figure 1C, 0.25 U of recombinant Dicer and 200 ng of recombinant PACT were used instead of the beads from immunoprecipitation. The reaction mixture was incubated at 37°C for 90 min. RNA was extracted from the reaction mixture by phenol extraction and analyzed on 15% denaturing polyacrylamide gel. Pre-let7a-1 was prepared from *in vitro* processing of cold pri-let7a-1 using Drosha-FLAG followed by gel purification. Pre-miR-30a and pre-let7a-1 were labeled at the 5' end using T4 polynucleotide kinase (TAKARA) and [ $\gamma$ - $^{32}$ P]ATP.

### RNAi and luciferase assay

HeLa cells cultured in six-well plates were cotransfected with 720 ng of pGL3-CMV firefly luciferase, 80 ng of pRL-CMV renilla luciferase, and 20 nM of siRNA against firefly luciferase, and 100 nM of siRNA against GFP, Dicer, PACT or TRBP. Lipofectamine 2000 (Invitrogen) was used for transfections, using the manufacturer's protocols. Transfections were performed in triplicate. Cells were lysed 48 h after transfection and assayed for luciferase activity using the Dual-Luciferase Reporter system (Promega) as described by the manufacturer. All firefly luciferase activities were normalized to renilla luciferase activities to correct for transfection efficiency. The target sequence of siDicer has been described previously (Lee *et al*, 2003). The target sequences of siPACT-1, siPACT-2, siTRBP, siExportin5, siAgo2, siGFP, and siLuciferase are 5'-GAACCAGCUUAAUCCUUAU-3' (siPACT-1), 5'-AGGAAUGCUGCUGAGAAAU-3' (siPACT-2), 5'-CACGUCAGGCUUACCUUGUAU-3' (siTRBP), 5'-CUCCAUUGGAGAAGGUGUA-3' (siExportin5), 5'-GCACGGAAGUCCAUUGUAU-3' (siAgo2), 5'-UGAAUUAGAUGGCGAUGUU-3' (siGFP), and 5'-CUUACGUGAGUACUUGCA-3' (siLuciferase), respectively. All siRNAs were manufactured by Samchully Pham Co. Ltd.

### Purification of recombinant proteins

To prepare recombinant PACT, cDNA was subcloned into pGEX-4T3 vector (Amersham), which has GST tag at the N-terminus, using *Bam*HI and *Xho*I sites. The PACT expression clone was transformed to the *E. coli* BL21-RIL strain. The expression and purification of recombinant PACT was conducted according to the manufacturer's protocol. After purification of GST-PACT, GST was removed from GST-PACT by treatment with 1 U/ml of Thrombin at 16°C for 16 h. TRBP protein was prepared similarly. Both recombinant proteins were used for *in vitro* assays and for immunization to raise polyclonal antibodies in rabbits.

### Northern blot analysis

To prepare the pri-miR-30a or pri-miR-124a inducible cell line, the insert containing pri-miR-30a genomic sequences, which was excised from pGEM-T-easy-pri-miR-30a (Lee et al, 2002) using *NotI* restriction enzyme, was subcloned into pTRE2-Hyg vector (Clontech) using *NotI* sites. Pri-miR-124a expression construct was cloned by PCR amplification of HeLa genomic DNA and inserted into the *NotI* site of the pTRE2-Hyg vector (Clontech). Tet-on HeLa cell line (Clontech) was transfected with the pri-miR-30a or pri-miR-124a inducible expression vector using Lipofectamine 2000 (Invitrogen). The selection of inducible cell line was conducted according to the manufacturer's protocol.

Pri-miR-30a or pri-miR-124a inducible cell line cultured in six-well plates was transfected with 100 nM of siRNAs using Lipofectamine 2000 (Invitrogen). In all, 3 µg/ml of doxycycline was added after 48 h. Then, total RNA was isolated using TRIzol reagent (Invitrogen) after 24 h. Total RNAs (25 µg) were resolved on 12.5% urea-polyacrylamide gels and transferred electronically to Zeta-probe membrane (BioRad). An oligonucleotide complementary to miR-30a-5p or miR-124a was end-labeled with [ $\gamma$ - $^{32}$ P]ATP and used as the probe for Northern.

### RT-PCR

In all, 5 µg of HeLa total RNA was used for the first-strand cDNA synthesis with SUPERScript II and oligo-dT primers (Invitrogen). To detect the expression level of mRNAs, the following primers

were used. For PACT, 5'-GGATCCATGTCAGAGCAGGCACC-3' (forward) and 5'-CTCGAGCTCCAGATTTACTTTCTTCTGC-3' (reverse); for TRBP, 5'-GGATCCATGAGTGAAGAGGAGCAAGGCTC-3' (forward) and 5'-CTCGAGTCAGGCAGTGAAGAGTCTGCTG-3' (reverse); for Dicer, 5'-AAGCTTATCCAGAGTGTTTGAGGGATAG-3' (forward) and 5'-CACAGTCAGGCTGATCAG-3' (reverse); for GAPDH, 5'-CCCATCACCATTCTCCAGGAGTGAAGAC-3' (forward) and 5'-CGCCC CACTTGATTTGGAGGGATCTCGCCTACCG-3' were used for PCR amplification.

### Supplementary data

Supplementary data are available at *The EMBO Journal* Online.

## Acknowledgements

We thank Drs Zissimos Mourelatos, Roe Hyun Seong, and Joo Yong Lee, and the members of our laboratory for their input and participation in helpful discussions. This work was supported by the Basic Research Program (R02-2004-000-10173-0) from the Ministry of Education and Human Resources Development and by the National Research Laboratory Program (M1050000010905J00 0010910), the SRC program of KOSEF (R11-2005-009-01003-0), and the Molecular and Cellular BioDiscovery Research Program (2004-01749) from the Ministry of Science and Technology.

## References

- Bartel DP (2004) MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell* **116**: 281–297
- Baulcombe D (2005) RNA silencing. *Trends Biochem Sci* **30**: 290–293
- Bernstein E, Caudy AA, Hammond SM, Hannon GJ (2001) Role for a bidentate ribonuclease in the initiation step of RNA interference. *Nature* **409**: 363–366
- Bernstein E, Kim SY, Carmell MA, Murchison EP, Alcorn H, Li MZ, Mills AA, Elledge SJ, Anderson KV, Hannon GJ (2003) Dicer is essential for mouse development. *Nat Genet* **35**: 215–217
- Bohnsack MT, Czaplinski K, Gorlich D (2004) Exportin 5 is a RanGTP-dependent dsRNA-binding protein that mediates nuclear export of pre-miRNAs. *RNA* **10**: 185–191
- Cai X, Hagedorn CH, Cullen BR (2004) Human microRNAs are processed from capped, polyadenylated transcripts that can also function as mRNAs. *RNA* **10**: 1957–1966
- Carmell MA, Xuan Z, Zhang MQ, Hannon GJ (2002) The Argonaute family: tentacles that reach into RNAi, developmental control, stem cell maintenance, and tumorigenesis. *Genes Dev* **16**: 2733–2742
- Chendrimada TP, Gregory RI, Kumaraswamy E, Norman J, Cooch N, Nishikura K, Shiekhattar R (2005) TRBP recruits the Dicer complex to Ago2 for microRNA processing and gene silencing. *Nature* **436**: 740–744
- Denli AM, Tops BB, Plasterk RH, Ketting RF, Hannon GJ (2004) Processing of primary microRNAs by the microprocessor complex. *Nature* **432**: 231–235
- Doi N, Zenno S, Ueda R, Ohki-Hamazaki H, Ui-Tei K, Saigo K (2003) Short-interfering-RNA-mediated gene silencing in mammalian cells requires dicer and eIF2C translation initiation factors. *Curr Biol* **13**: 41–46
- Filipowicz W (2005) RNAi: the nuts and bolts of the RISC machine. *Cell* **122**: 17–20
- Forstemann K, Tomari Y, Du T, Vagin VV, Denli AM, Bratu DP, Klattenhoff C, Theurkauf WE, Zamore PD (2005) Normal microRNA maturation and germ-line stem cell maintenance requires Loquacious, a double-stranded RNA-binding domain protein. *PLoS Biol* **3**: e236
- Gregory RI, Chendrimada TP, Cooch N, Shiekhattar R (2005) Human RISC couples MicroRNA biogenesis and posttranscriptional gene silencing. *Cell* **123**: 631–640
- Gregory RI, Yan KP, Amuthan G, Chendrimada T, Doratotaj B, Cooch N, Shiekhattar R (2004) The microprocessor complex mediates the genesis of microRNAs. *Nature* **432**: 235–240
- Grishok A, Pasquinelli AE, Conte D, Li N, Parrish S, Ha I, Baillie DL, Fire A, Ruvkun G, Mello CC (2001) Genes and mechanisms related to RNA interference regulate expression of the small temporal RNAs that control *C. elegans* developmental timing. *Cell* **106**: 23–34
- Gupta V, Huang X, Patel RC (2003) The carboxy-terminal, M3 motifs of PACT and TRBP have opposite effects on PKR activity. *Virology* **315**: 283–291
- Haase AD, Jaskiewicz L, Zhang H, Laine S, Sack R, Gatignol A, Filipowicz W (2005) TRBP, a regulator of cellular PKR and HIV-1 virus expression, interacts with Dicer and functions in RNA silencing. *EMBO Rep* **6**: 961–967
- Hammond SM, Boettcher S, Caudy AA, Kobayashi R, Hannon GJ (2001) Argonaute2, a link between genetic and biochemical analyses of RNAi. *Science* **293**: 1146–1150
- Han J, Lee Y, Yeom KH, Kim YK, Jin H, Kim VN (2004) The Drosha-DGCR8 complex in primary microRNA processing. *Genes Dev* **18**: 3016–3027
- Hutvagner G, McLachlan J, Pasquinelli AE, Balint E, Tuschl T, Zamore PD (2001) A cellular function for the RNA-interference enzyme Dicer in the maturation of the let-7 small temporal RNA. *Science* **293**: 834–838
- Jiang F, Ye X, Liu X, Fincher L, McKearin D, Liu Q (2005) Dicer-1 and R3D1-L catalyze microRNA maturation in *Drosophila*. *Genes Dev* **19**: 1674–1679
- Ketting RF, Fischer SE, Bernstein E, Sijen T, Hannon GJ, Plasterk RH (2001) Dicer functions in RNA interference and in synthesis of small RNA involved in developmental timing in *C. elegans*. *Genes Dev* **15**: 2654–2659
- Kim VN (2005a) MicroRNA biogenesis: coordinated cropping and dicing. *Nat Rev Mol Cell Biol* **6**: 376–385
- Kim VN (2005b) Small RNAs: classification, biogenesis, and function. *Mol Cells* **19**: 1–15
- Landthaler M, Yalcin A, Tuschl T (2004) The human DiGeorge syndrome critical region gene 8 and its *D. melanogaster* homolog are required for miRNA biogenesis. *Curr Biol* **14**: 2162–2167
- Lee Y, Ahn C, Han J, Choi H, Kim J, Yim J, Lee J, Provost P, Radmark O, Kim S, Kim VN (2003) The nuclear RNase III Drosha initiates microRNA processing. *Nature* **425**: 415–419
- Lee Y, Jeon K, Lee JT, Kim S, Kim VN (2002) MicroRNA maturation: stepwise processing and subcellular localization. *EMBO J* **21**: 4663–4670
- Lee Y, Kim M, Han J, Yeom KH, Lee S, Baek SH, Kim VN (2004a) MicroRNA genes are transcribed by RNA polymerase II. *EMBO J* **23**: 4051–4060
- Lee YS, Nakahara K, Pham JW, Kim K, He Z, Sontheimer EJ, Carthew RW (2004b) Distinct roles for *Drosophila* Dicer-1 and Dicer-2 in the siRNA/miRNA silencing pathways. *Cell* **117**: 69–81

- Lingel A, Simon B, Izaurralde E, Sattler M (2004) Nucleic acid 3'-end recognition by the Argonaute2 PAZ domain. *Nat Struct Mol Biol* **11**: 576–577
- Liu J, Carmell MA, Rivas FV, Marsden CG, Thomson JM, Song JJ, Hammond SM, Joshua-Tor L, Hannon GJ (2004) Argonaute2 is the catalytic engine of mammalian RNAi. *Science* **305**: 1437–1441
- Liu Q, Rand TA, Kalidas S, Du F, Kim HE, Smith DP, Wang X (2003) R2D2, a bridge between the initiation and effector steps of the *Drosophila* RNAi pathway. *Science* **301**: 1921–1925
- Lund E, Guttinger S, Calado A, Dahlberg JE, Kutay U (2004) Nuclear export of microRNA precursors. *Science* **303**: 95–98
- Ma JB, Ye K, Patel DJ (2004) Structural basis for overhang-specific small interfering RNA recognition by the PAZ domain. *Nature* **429**: 318–322
- Ma JB, Yuan YR, Meister G, Pei Y, Tuschl T, Patel DJ (2005) Structural basis for 5'-end-specific recognition of guide RNA by the *A. fulgidus* Piwi protein. *Nature* **434**: 666–670
- Maniatakis E, Mourelatos Z (2005) A human, ATP-independent, RISC assembly machine fueled by pre-miRNA. *Genes Dev* **19**: 2979–2990
- Meister G, Landthaler M, Patkaniowska A, Dorsett Y, Teng G, Tuschl T (2004) Human Argonaute2 mediates RNA cleavage targeted by miRNAs and siRNAs. *Mol Cell* **15**: 185–197
- Meister G, Landthaler M, Peters L, Chen PY, Urlaub H, Luhrmann R, Tuschl T (2005) Identification of novel Argonaute-associated proteins. *Curr Biol* **15**: 2149–2155
- Parker JS, Roe SM, Barford D (2005) Structural insights into mRNA recognition from a PIWI domain-siRNA guide complex. *Nature* **434**: 663–666
- Patel RC, Sen GC (1998) PACT, a protein activator of the interferon-induced protein kinase, PKR. *EMBO J* **17**: 4379–4390
- Peters GA, Hartmann R, Qin J, Sen GC (2001) Modular structure of PACT: distinct domains for binding and activating PKR. *Mol Cell Biol* **21**: 1908–1920
- Pham JW, Pellino JL, Lee YS, Carthew RW, Sontheimer EJ (2004) A Dicer-2-dependent 80s complex cleaves targeted mRNAs during RNAi in *Drosophila*. *Cell* **117**: 83–94
- Pham JW, Sontheimer EJ (2005) Molecular requirements for RNA-induced silencing complex assembly in the *Drosophila* RNA interference pathway. *J Biol Chem* **280**: 39278–39283
- Rivas FV, Tolia NH, Song JJ, Aragon JP, Liu J, Hannon GJ, Joshua-Tor L (2005) Purified Argonaute2 and an siRNA form recombinant human RISC. *Nat Struct Mol Biol* **12**: 340–349
- Saito K, Ishizuka A, Siomi H, Siomi MC (2005) Processing of pre-microRNAs by the Dicer-1-Loquacious complex in *Drosophila* cells. *PLoS Biol* **3**: e235
- Saunders LR, Perkins DJ, Balachandran S, Michaels R, Ford R, Mayeda A, Barber GN (2001) Characterization of two evolutionarily conserved, alternatively spliced nuclear phosphoproteins, NFAR-1 and -2, that function in mRNA processing and interact with the double-stranded RNA-dependent protein kinase, PKR. *J Biol Chem* **276**: 32300–32312
- Song JJ, Liu J, Tolia NH, Schneiderman J, Smith SK, Martienssen RA, Hannon GJ, Joshua-Tor L (2003) The crystal structure of the Argonaute2 PAZ domain reveals an RNA binding motif in RNAi effector complexes. *Nat Struct Biol* **10**: 1026–1032
- Song JJ, Smith SK, Hannon GJ, Joshua-Tor L (2004) Crystal structure of Argonaute and its implications for RISC slicer activity. *Science* **305**: 1434–1437
- Sontheimer EJ (2005) Assembly and function of RNA silencing complexes. *Nat Rev Mol Cell Biol* **6**: 127–138
- Sontheimer EJ, Carthew RW (2004) Molecular biology. Argonaute journeys into the heart of RISC. *Science* **305**: 1409–1410
- Taylor SS, Haste NM, Ghosh G (2005) PKR and eIF2alpha: integration of kinase dimerization, activation, and substrate docking. *Cell* **122**: 823–825
- Tomari Y, Du T, Haley B, Schwarz DS, Bennett R, Cook HA, Koppetsch BS, Theurkauf WE, Zamore PD (2004a) RISC assembly defects in the *Drosophila* RNAi mutant armitage. *Cell* **116**: 831–841
- Tomari Y, Matranga C, Haley B, Martinez N, Zamore PD (2004b) A protein sensor for siRNA asymmetry. *Science* **306**: 1377–1380
- Tomari Y, Zamore PD (2005) Perspective: machines for RNAi. *Genes Dev* **19**: 517–529
- Xu Z, Williams BR (2000) The B56alpha regulatory subunit of protein phosphatase 2A is a target for regulation by double-stranded RNA-dependent protein kinase PKR. *Mol Cell Biol* **20**: 5285–5299
- Yan KS, Yan S, Farooq A, Han A, Zeng L, Zhou MM (2003) Structure and conserved RNA binding of the PAZ domain. *Nature* **426**: 468–474
- Yi R, Qin Y, Macara IG, Cullen BR (2003) Exportin-5 mediates the nuclear export of pre-microRNAs and short hairpin RNAs. *Genes Dev* **17**: 3011–3016
- Yuan YR, Pei Y, Ma JB, Kuryavyy V, Zhadina M, Meister G, Chen HY, Dauter Z, Tuschl T, Patel DJ (2005) Crystal structure of *A. aeolicus* argonaute, a site-specific DNA-guided endoribonuclease, provides insights into RISC-mediated mRNA cleavage. *Mol Cell* **19**: 405–419
- Zamore PD, Haley B (2005) Ribo-gnome: the big world of small RNAs. *Science* **309**: 1519–1524
- Zhong J, Peters AH, Lee K, Braun RE (1999) A double-stranded RNA binding protein required for activation of repressed messages in mammalian germ cells. *Nat Genet* **22**: 171–174